

GEPHE SUMMARY

<p>WntA (https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=~WntA^#gephebase-summary-title)</p> <p>Published</p>	<p>Gephebase Gene</p> <p>Entry Status</p>	<p>GP00001207</p> <p>Martin</p>	<p>GepheID</p> <p>Main curator</p>
--	---	---------------------------------	------------------------------------

PHENOTYPIC CHANGE

<p>Morphology (https://www.gephebase.org/search-criteria?/and+Trait+Category=~Morphology^#gephebase-summary-title)</p> <p>Coloration (wing, Mullerian mimicry) (https://www.gephebase.org/search-criteria?/and+Trait=~Coloration+(wing,+Mullerian+mimicry)^#gephebase-summary-title)</p> <p>Heliconius ismenius bouletti</p> <p>Heliconius ismenius telchinia - differs notably by coverage of melanic and WntA-positive patches</p> <p>Unknown</p> <p>Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=~Intraspecific^#gephebase-summary-title)</p>	<p>Trait Category</p> <p>Trait</p> <p>Trait State in Taxon A</p> <p>Trait State in Taxon B</p> <p>Ancestral State</p> <p>Taxonomic Status</p>	<p>Taxon A</p> <p>Latin Name</p> <p>Heliconius ismenius (https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Heliconius+ismenius^#gephebase-summary-title)</p> <p>-</p> <p>Heliconius ismenius Latreille, 1817</p> <p>species</p> <p>cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphimesnoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Dityisia; Obtectomera; Papilionoidea; Nymphalidae; Heliconiinae; Heliconiini; Heliconius</p> <p>Heliconius () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=33416)</p> <p>33427 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=33427)</p> <p>is Taxon A an Infrasppecies?</p> <p>Yes</p> <p>Heliconius ismenius bouletti</p>	<p>Latin Name</p> <p>Common Name</p> <p>Synonyms</p> <p>Rank</p> <p>Lineage</p> <p>Parent</p> <p>NCBI Taxonomy ID</p> <p>is Taxon B an Infrasppecies?</p> <p>Taxon B Description</p>	<p>Taxon B</p> <p>Latin Name</p> <p>Heliconius ismenius (https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Heliconius+ismenius^#gephebase-summary-title)</p> <p>-</p> <p>Heliconius ismenius Latreille, 1817</p> <p>species</p> <p>cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphimesnoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Dityisia; Obtectomera; Papilionoidea; Nymphalidae; Heliconiinae; Heliconiini; Heliconius</p> <p>Heliconius () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=33416)</p> <p>33427 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=33427)</p> <p>is Taxon B an Infrasppecies?</p> <p>Yes</p> <p>Heliconius ismenius telchinia - differs notably by coverage of melanic and WntA-positive patches</p>
---	---	--	--	--

GENOTYPIC CHANGE

<p>WntA</p> <p>-</p> <p>-</p> <p>Belongs to the Wnt family.</p> <p>GO:0005102 : signaling receptor binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005102)</p> <p>GO:0007275 : multicellular organism development</p>	<p>Generic Gene Name</p> <p>Synonyms</p> <p>String</p> <p>Sequence Similarities</p> <p>GO - Molecular Function</p> <p>GO - Biological Process</p>	<p>A0A077DF90 (http://www.uniprot.org/uniprot/A0A077DF90)</p> <p>()</p>	<p>UniProtKB Vanessa cardui</p> <p>GenebankID or UniProtKB</p>
---	---	---	--

(<https://www.ebi.ac.uk/QuickGO/term/GO:0007275>)

GO:0016055 : Wnt signaling pathway

(<https://www.ebi.ac.uk/QuickGO/term/GO:0016055>)

GO - Cellular Component

GO:0005576 : extracellular region (<https://www.ebi.ac.uk/QuickGO/term/GO:0005576>)

Presumptive Null

No ([https://www.gephebase.org/search-criteria?/and+Presumptive Null="+No^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Presumptive+Null=))

Molecular Type

Cis-regulatory ([https://www.gephebase.org/search-criteria?/and+Molecular Type="+Cis-regulatory^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Molecular+Type=))

Aberration Type

Unknown ([https://www.gephebase.org/search-criteria?/and+Aberration Type="+Unknown^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Aberration+Type=))

Molecular Details of the Mutation

unknown

Experimental Evidence

Linkage Mapping ([https://www.gephebase.org/search-criteria?/and+Experimental Evidence="+Linkage Mapping^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=))

Main Reference

Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. (2012) (<https://pubmed.ncbi.nlm.nih.gov/22802635>)

Authors

Martin A; Papa R; Nadeau NJ; Hill RI; Counterman BA; Halder G; Jiggins CD; Kronforst MR; Long AD; McMillan WO; Reed RD

Abstract

Although animals display a rich variety of shapes and patterns, the genetic changes that explain how complex forms arise are still unclear. Here we take advantage of the extensive diversity of *Heliconius* butterflies to identify a gene that causes adaptive variation of black wing patterns within and between species. Linkage mapping in two species groups, gene-expression analysis in seven species, and pharmacological treatments all indicate that cis-regulatory evolution of the WntA ligand underpins discrete changes in color pattern features across the *Heliconius* genus. These results illustrate how the direct modulation of morphogen sources can generate a wide array of unique morphologies, thus providing a link between natural genetic variation, pattern formation, and adaptation.

Additional References

RELATED GEPHE

No matches found.

Related Genes

No matches found.

Related Haplotypes

EXTERNAL LINKS

COMMENTS