

RDO5 REDUCED DORMANCY5 (https://www.gephebase.org/search-criteria?/and+Gene=Gephebase=RDO5 REDUCED DORMANCY5^#gephebase-summary-title)	Gephebase Gene	GP00001248	GepheID
Published	Entry Status	Arnoult	Main curator

	Trait Category
Physiology (https://www.gephebase.org/search-criteria?/and+Trait+Category=~Physiology^#gephebase-summary-title)	
	Trait
Seed dormancy (https://www.gephebase.org/search-criteria?/and+Trait=~Seed+dormancy^#gephebase-summary-title)	
	Trait State in Taxon A
Arabidopsis thaliana- Ler0	
	Trait State in Taxon B
Arabidopsis thaliana Love-1	
	Ancestral State
Taxon A	
	Taxonomic Status
Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=~Intraspecific^#gephebase-summary-title)	

GENOTYPIC CHANGE		
At4g11040	Generic Gene Name	UniProtKB Arabidopsis thaliana
	Synonyms	GenebankID or UniProtKB
RDO5; REDUCED DORMANCY 5; T22B4.20; T22B4_20; At4g11040; F8M12.14	String	
3702.AT4G11040.1 (http://string-db.org/newstring.cgi/show_network_section.pl?identifier= 3702.AT4G11040.1)	Sequence Similarities	
Belongs to the PP2C family.	GO - Molecular Function	
	GO:0046872 : metal ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0046872)	
	GO:0004721 : phosphoprotein phosphatase activity (https://www.ebi.ac.uk/QuickGO/term/GO:0004721)	

GO:0004722 : protein serine/threonine phosphatase activity
(<https://www.ebi.ac.uk/QuickGO/term/GO:0004722>)
GO:0001691 : pseudophosphatase activity
(<https://www.ebi.ac.uk/QuickGO/term/GO:0001691>)

GO - Biological Process

GO:0010162 : seed dormancy process
(<https://www.ebi.ac.uk/QuickGO/term/GO:0010162>)
GO:1902040 : positive regulation of seed dormancy process
(<https://www.ebi.ac.uk/QuickGO/term/GO:1902040>)

GO - Cellular Component

GO:0005634 : nucleus (<https://www.ebi.ac.uk/QuickGO/term/GO:0005634>)

Presumptive Null

Yes (<https://www.gephebase.org/search-criteria?/and+Presumptive Null=~Yes~#gephebase-summary-title>)

Molecular Type

Coding (<https://www.gephebase.org/search-criteria?/and+Molecular Type=~Coding~#gephebase-summary-title>)

Aberration Type

Deletion (<https://www.gephebase.org/search-criteria?/and+Aberration Type=~Deletion~#gephebase-summary-title>)

Deletion Size

10-99 bp

Molecular Details of the Mutation

-23bp at position 1371

Experimental Evidence

Linkage Mapping (<https://www.gephebase.org/search-criteria?/and+Experimental Evidence=~Linkage Mapping~#gephebase-summary-title>)

Main Reference

Sequence Polymorphisms at the REDUCED DORMANCY5 Pseudophosphatase Underlie Natural Variation in Arabidopsis Dormancy. (2016) (<https://pubmed.ncbi.nlm.nih.gov/27288362>)

Authors

Xiang Y; Song B; NÄ©e G; Kramer K; Finkemeier I; Soppe WJ

Abstract

Seed dormancy controls the timing of germination, which regulates the adaptation of plants to their environment and influences agricultural production. The time of germination is under strong natural selection and shows variation within species due to local adaptation. The identification of genes underlying dormancy quantitative trait loci is a major scientific challenge, which is relevant for agricultural and ecological goals. In this study, we describe the identification of the DELAY OF GERMINATION18 (DOG18) quantitative trait locus, which was identified as a factor in natural variation for seed dormancy in Arabidopsis (*Arabidopsis thaliana*). DOG18 encodes a member of the clade A of the type 2C protein phosphatases family, which we previously identified as the REDUCED DORMANCY5 (RDO5) gene. DOG18/RDO5 shows a relatively high frequency of loss-of-function alleles in natural accessions restricted to northwestern Europe. The loss of dormancy in these loss-of-function alleles can be compensated for by genetic factors like DOG1 and DOG6, and by environmental factors such as low temperature. RDO5 does not have detectable phosphatase activity. Analysis of the phosphoproteome in dry and imbibed seeds revealed a general decrease in protein phosphorylation during seed imbibition that is enhanced in the rdo5 mutant. We conclude that RDO5 acts as a pseudophosphatase that inhibits dephosphorylation during seed imbibition.

Â© 2016 American Society of Plant Biologists. All Rights Reserved.

Additional References

Natural variation for seed dormancy in Arabidopsis is regulated by additive genetic and molecular pathways. (2010) (<https://pubmed.ncbi.nlm.nih.gov/20145108>)

RELATED GEPHE

Related Genes

2 (D14 (KAI2 paralogue), DOG1 (DELAY OF GERMINATION 1)) (<https://www.gephebase.org/search-criteria?/or+Taxon ID=~3702~/and+Trait=Seed dormancy/and+groupHaplotypes=true#gephebase-summary-title>)

Related Haplotypes

15 (<https://www.gephebase.org/search-criteria?/or+Gene Gephebase=~RDO5 REDUCED DORMANCY5~/and+Taxon ID=~3702~/or+Gene Gephebase=~RDO5 REDUCED DORMANCY5~/and+Taxon ID=~3702~#gephebase-summary-title>)

EXTERNAL LINKS

COMMENTS

predicted phenotypic effect