

GEPHE SUMMARY

tyrosinase (TYR) (https://www.gephebase.org/search-criteria?/and+Gene) Gephebase= [^] tyrosinase (TYR) [^] #gephebase-summary-title)	Gephebase Gene	GP00001347	GepheID
Published	Entry Status	Prigent	Main curator

PHENOTYPIC CHANGE

Morphology (https://www.gephebase.org/search-criteria?/and+Trait) Category= [^] Morphology [^] #gephebase-summary-title)	Trait Category		
Coloration (skin; coat) (https://www.gephebase.org/search-criteria?/and+Trait) (skin; coat) [^] #gephebase-summary-title)	Trait		
Donkey ; wild type	Trait State in Taxon A		
Donkey- Asinara white ; albino	Trait State in Taxon B		
Taxon A	Ancestral State		
Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic) Status= [^] Intraspecific [^] #gephebase-summary-title)	Taxonomic Status		
	Taxon A	Taxon B	
Equus asinus (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms= [^] Equus asinus [^] #gephebase-summary-title)	Latin Name	Equus asinus (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms= [^] Equus asinus [^] #gephebase-summary-title)	Latin Name
ass	Common Name	ass	Common Name
ass; African ass; African wild ass; Somali wild ass; domestic ass; donkey	Synonyms	ass; African ass; African wild ass; Somali wild ass; domestic ass; donkey	Synonyms
species	Rank	species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Sarcopterygii; Dipnotetrapodomorpha; Tetrapoda; Amniota; Mammalia; Theria; Eutheria; Boreoeutheria; Laurasiatheria; Perissodactyla; Equidae; Equus; Asinus	Lineage	cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Sarcopterygii; Dipnotetrapodomorpha; Tetrapoda; Amniota; Mammalia; Theria; Eutheria; Boreoeutheria; Laurasiatheria; Perissodactyla; Equidae; Equus; Asinus	Lineage
Asinus () - (Rank: subgenus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=35508)	Parent	Asinus () - (Rank: subgenus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=35508)	Parent
9793 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=9793)	NCBI Taxonomy ID	9793 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=9793)	NCBI Taxonomy ID
No	is Taxon A an Intraspecies?	Yes	is Taxon B an Intraspecies?
		Donkey- Asinara white ; albino	Taxon B Description

GENOTYPIC CHANGE

Tyr	Generic Gene Name	P11344 (http://www.uniprot.org/uniprot/P11344)	UniProtKB Mus musculus
c; Oca1; skc35; albino	Synonyms	()	GenebankID or UniProtKB
10090.ENSMUSP00000004770 (http://string-db.org/newstring.cgi/show_network_section.pl?identifier=10090.ENSMUSP00000004770)	String		
Belongs to the tyrosinase family.	Sequence Similarities		
GO:0042803 : protein homodimerization activity (https://www.ebi.ac.uk/QuickGO/term/GO:0042803)	GO - Molecular Function		
GO:0046982 : protein heterodimerization activity (https://www.ebi.ac.uk/QuickGO/term/GO:0046982)			
GO:0005507 : copper ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005507)			

GO:0004503 : monophenol monooxygenase activity
(<https://www.ebi.ac.uk/QuickGO/term/GO:0004503>)

GO - Biological Process

GO:0042438 : melanin biosynthetic process

(<https://www.ebi.ac.uk/QuickGO/term/GO:0042438>)

GO:0043473 : pigmentation (<https://www.ebi.ac.uk/QuickGO/term/GO:0043473>)

GO:0008283 : cell proliferation (<https://www.ebi.ac.uk/QuickGO/term/GO:0008283>)

GO:0033280 : response to vitamin D (<https://www.ebi.ac.uk/QuickGO/term/GO:0033280>)

GO:0051591 : response to cAMP (<https://www.ebi.ac.uk/QuickGO/term/GO:0051591>)

GO:0009411 : response to UV (<https://www.ebi.ac.uk/QuickGO/term/GO:0009411>)

GO:0048538 : thymus development (<https://www.ebi.ac.uk/QuickGO/term/GO:0048538>)

GO - Cellular Component

GO:0016021 : integral component of membrane

(<https://www.ebi.ac.uk/QuickGO/term/GO:0016021>)

GO:0005737 : cytoplasm (<https://www.ebi.ac.uk/QuickGO/term/GO:0005737>)

GO:0005829 : cytosol (<https://www.ebi.ac.uk/QuickGO/term/GO:0005829>)

GO:0005634 : nucleus (<https://www.ebi.ac.uk/QuickGO/term/GO:0005634>)

GO:0043231 : intracellular membrane-bounded organelle

(<https://www.ebi.ac.uk/QuickGO/term/GO:0043231>)

GO:0048471 : perinuclear region of cytoplasm

(<https://www.ebi.ac.uk/QuickGO/term/GO:0048471>)

GO:0042470 : melanosome (<https://www.ebi.ac.uk/QuickGO/term/GO:0042470>)

GO:0033162 : melanosome membrane

(<https://www.ebi.ac.uk/QuickGO/term/GO:0033162>)

Presumptive Null

No (<https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title>)

Molecular Type

Coding (<https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title>)

Aberration Type

SNP (<https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title>)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

c.604C>G p.His202Asp

Experimental Evidence

Candidate Gene (<https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title>)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	His	Asp	202

Main Reference

The albinism of the feral Asinara white donkeys (*Equus asinus*) is determined by a missense mutation in a highly conserved position of the tyrosinase (TYR) gene deduced protein. (2016)
(<https://pubmed.ncbi.nlm.nih.gov/26763160>)

Authors

Utzeri VJ; Bertolini F; Ribani A; Schiavo G; Dall'Olio S; Fontanesi L

Abstract

A feral donkey population (*Equus asinus*), living in the Asinara National Park (an island north-west of Sardinia, Italy), includes a unique white albino donkey subpopulation or colour morph that is a major attraction of this park. Disrupting mutations in the tyrosinase (TYR) gene are known to cause recessive albinisms in humans (oculocutaneous albinism Type 1; OCA1) and other species. In this study, we analysed the donkey TYR gene as a strong candidate to identify the causative mutation of the albinism of these donkeys. The TYR gene was sequenced from 13 donkeys (seven Asinara white albino and six coloured animals). Seven single nucleotide polymorphisms were identified. A missense mutation (c.604C>G; p.His202Asp) in a highly conserved amino acid position (even across kingdoms), which disrupts the first copper-binding site (CuA) of functional protein, was identified in the homozygous condition (G/G or D/D) in all Asinara white albino donkeys and in the albino son of a trio (the grey parents had genotype C/G or H/D), supporting the recessive mode of inheritance of this mutation. Genotyping 82 donkeys confirmed that Asinara albino donkeys had genotype G/G whereas all other coloured donkeys had genotype C/C or C/G. Across-population association between the c.604C>G genotypes and the albino coat colour was highly significant (P = 6.17E-18). The identification of the causative mutation of the albinism in the Asinara white donkeys might open new perspectives to study the dynamics of this putative deleterious allele in a feral population and to manage this interesting animal genetic resource.

Â© 2015 Stichting International Foundation for Animal Genetics.

Additional References

RELATED GEPHE

Related Genes

3 (Agouti (ASIP), Kit (type III receptor protein-tyrosine kinase), MC1R) (<https://www.gephebase.org/search-criteria?/or+Taxon ID=^9793^/and+Trait=Coloration/and+groupHaplotypes=true#gephebase-summary-title>)

Related Haplotypes

No matches found.

EXTERNAL LINKS

COMMENTS

recessive ; @Parallelism <https://omia.org/OMIA000202/9793/>