

GEPHE SUMMARY

	Gephebase Gene		GepheID
Heading Date 1 (HD1) (https://www.gephebase.org/search-criteria?/and+Gene)		GP00001408	
Gephebase="Heading Date 1 (HD1)"#gephebase-summary-title)			Main curator
	Entry Status	Prigent	
Published			

PHENOTYPIC CHANGE

	Trait Category		
Physiology (https://www.gephebase.org/search-criteria?/and+Trait)			
Category="Physiology"#gephebase-summary-title)	Trait		
Flowering time (<a flowering"="" href="https://www.gephebase.org/search-criteria?/and+Trait=">https://www.gephebase.org/search-criteria?/and+Trait="Flowering)			
time"#gephebase-summary-title)	Trait State in Taxon A		
early flowering Domesticated sorghum line TX623			
	Trait State in Taxon B		
Late flowering wild sorghum & Domesticated sorghum from Ethiopia Yemen Germany & China			
	Ancestral State		
Taxon A			
	Taxonomic Status		
Domesticated (https://www.gephebase.org/search-criteria?/and+Taxonomic)			
Status="Domesticated"#gephebase-summary-title)			
	Taxon A	Taxon B	
	Latin Name		Latin Name
Sorghum bicolor		Sorghum virgatum	
(https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="Sorghum)		(https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="Sorghum)	
bicolor"#gephebase-summary-title)	Common Name		Common Name
sorghum		-	
	Synonyms		Synonyms
Andropogon sorghum; Sorghum bicolor subsp. bicolor; Sorghum nervosum; Sorghum saccharatum; Sorghum vulgare; sorghum; broomcorn; milo; Andropogon sorghum (L.) Brot.; Sorghum bicolor (L.) Moench; Sorghum nervosum Besser ex Schul.; Sorghum saccharatum (L.) Moench; Sorghum vulgare Pers.; Sorghum bicolor milo; Sorghum_bicolor		Sorghum bicolor var. virgatum; Sorghum bicolor var. virgatum (Hack.) de Wet & Huckabay, nom. inval.; Sorghum virgatum (Hack.) Stapf	
	Rank		Rank
species		species	
	Lineage		Lineage
cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliopsida; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; PACMAD clade; Panicoideae; Andropogonodae; Andropogoneae; Sorghinae; Sorghum		cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliophyta; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; PACMAD clade; Panicoideae; Andropogonodae; Andropogoneae; Sorghinae; Sorghum	
	Parent		Parent
Sorghum () - (Rank: genus)		Sorghum () - (Rank: genus)	
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 4557)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 4557)	
4558	NCBI Taxonomy ID	1428165	NCBI Taxonomy ID
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 4558)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 1428165)	
	is Taxon A an Intraspecies?		is Taxon B an Intraspecies?
Yes		Yes	
	Taxon A Description		Taxon B Description
early flowering Domesticated sorghum line TX623		Late flowering wild sorghum & Domesticated sorghum from Ethiopia Yemen Germany & China	

GENOTYPIC CHANGE

	Generic Gene Name		UniProtKB Arabidopsis thaliana
CO		Q39057 (http://www.uniprot.org/uniprot/Q39057)	
	Synonyms		GenebankID or UniProtKB
B-box domain protein 1; BBX1; CONSTANS; F14F8.220; F14F8_220; FG; At5g15840		()	
	String		
3702.AT5G15840.1			
(http://string-db.org/newstring.cgi/show_network_section.pl?identifier= 3702.AT5G15840.1)			
	Sequence Similarities		
Belongs to the CONSTANS family.			
	GO - Molecular Function		
GO:0003700 : DNA-binding transcription factor activity			

(<https://www.ebi.ac.uk/QuickGO/term/GO:0003700>)
 GO:0008270 : zinc ion binding (<https://www.ebi.ac.uk/QuickGO/term/GO:0008270>)
 GO:0003677 : DNA binding (<https://www.ebi.ac.uk/QuickGO/term/GO:0003677>)
 GO - Biological Process
 GO:0030154 : cell differentiation (<https://www.ebi.ac.uk/QuickGO/term/GO:0030154>)
 GO:0009908 : flower development (<https://www.ebi.ac.uk/QuickGO/term/GO:0009908>)
 GO:0009909 : regulation of flower development
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0009909>)
 GO:0010218 : response to far red light
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0010218>)
 GO:0010018 : far-red light signaling pathway
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0010018>)

GO - Cellular Component

GO:0005634 : nucleus (<https://www.ebi.ac.uk/QuickGO/term/GO:0005634>)

Presumptive Null

Yes ([https://www.gephebase.org/search-criteria?/and+Presumptive Null=~Yes~#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Presumptive+Null=~Yes~#gephebase-summary-title))

Molecular Type

Coding ([https://www.gephebase.org/search-criteria?/and+Molecular Type=~Coding~#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Molecular+Type=~Coding~#gephebase-summary-title))

Aberration Type

Deletion ([https://www.gephebase.org/search-criteria?/and+Aberration Type=~Deletion~#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Aberration+Type=~Deletion~#gephebase-summary-title))

Deletion Size

1-9 bp

Molecular Details of the Mutation

5bp deletion in the coding sequence leading to gene frameshift

Experimental Evidence

Association Mapping ([https://www.gephebase.org/search-criteria?/and+Experimental Evidence=~Association Mapping~#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=~Association+Mapping~#gephebase-summary-title))

Main Reference

Parallel Domestication of the Heading Date 1 Gene in Cereals. (2015) (<https://pubmed.ncbi.nlm.nih.gov/26116860>)

Authors

Liu H; Liu H; Zhou L; Zhang Z; Zhang X; Wang M; Li H; Lin Z

Abstract

Flowering time is one of the key determinants of crop adaptation to local environments during domestication. However, the genetic basis underlying flowering time is yet to be elucidated in most cereals. Although staple cereals, such as rice, maize, wheat, barley, and sorghum, have spread and adapted to a wide range of ecological environments during domestication, it is yet to be determined whether they have a common genetic basis for flowering time. In this study, we show, through map-based cloning, that flowering time in sorghum is controlled by a major quantitative trait locus (QTL) Heading Date 1 (HD1), located on chromosome 10. The causal gene encodes the CONSTANS gene family which contains a CCT domain. A 5-bp deletion of a minor allele present in the coding sequence leads to a gene frameshift that delays flowering in sorghum. In contrast, in foxtail millet, association mapping of HD1 showed a common causal site with a splicing variant from "GT" to "AT" that was highly correlated with flowering time. In addition, the rice HD1 gene is known to harbor several causal variants controlling flowering time. These data indicate that the major flowering time QTL HD1 was under parallel domestication in sorghum, foxtail millet, and rice. The pattern of common mixed minor, or even rare, causal alleles in HD1 across different species may be representative of the genetic basis of the domestication syndrome. Furthermore, large DNA sequence analysis of HD1 revealed multiple origins for domesticated sorghum and a single origin for domesticated foxtail millet.

© The Author 2015. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

Additional References

RELATED GEPHE

Related Genes

1 (PRR37 pseudoresponse regulator protein 37) ([https://www.gephebase.org/search-criteria?/or+Taxon ID=~4558~/and+Trait=Flowering time/or+Taxon ID=~1428165~/and+Trait=Flowering time/and+groupHaplotypes=true#gephebase-summary-title](https://www.gephebase.org/search-criteria?/or+Taxon+ID=~4558~/and+Trait=Flowering+time/or+Taxon+ID=~1428165~/and+Trait=Flowering+time/and+groupHaplotypes=true#gephebase-summary-title))

Related Haplotypes

No matches found.

EXTERNAL LINKS

COMMENTS

it is unclear in which allele is the mutation and paper is confusing whether wild type is early or late flowering