

GEPHE SUMMARY

	Gephebase Gene		GepheID
Early flowering1 (EL1) (https://www.gephebase.org/search-criteria?/and+Gene)		GP00001636	
Gephebase="Early flowering1 (EL1)"#gephebase-summary-title)			Main curator
Published	Entry Status	Prigent	

PHENOTYPIC CHANGE

	Trait Category		
Physiology (https://www.gephebase.org/search-criteria?/and+Trait)			
Category="Physiology"#gephebase-summary-title)	Trait		
Flowering time (heading date) (<a (heading="" date)"#gephebase-summary-title"="" flowering="" href="https://www.gephebase.org/search-criteria?/and+Trait=" time="">https://www.gephebase.org/search-criteria?/and+Trait="Flowering time (heading date)"#gephebase-summary-title)			
	Trait State in Taxon A		
Middle-late-flowering Tongil-type (japonica/indica hybrid) rice Milyang23 (116 days)			
	Trait State in Taxon B		
Early heading japonica rice H143 (78 days) and H75 (Hokkaido; Japan)			
	Ancestral State		
Taxon A			
	Taxonomic Status		
Domesticated (https://www.gephebase.org/search-criteria?/and+Taxonomic)			
Status="Domesticated"#gephebase-summary-title)			
Taxon A		Taxon B	
	Latin Name		Latin Name
Oryza sativa		Oryza sativa Japonica Group	
(https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="Oryza sativa"#gephebase-summary-title)		(<a group"#gephebase-summary-title"="" href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=" japonica="" oryza="" sativa="">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="Oryza sativa Japonica Group"#gephebase-summary-title)	
	Common Name		Common Name
rice		Japanese rice	
	Synonyms		Synonyms
rice; red rice; Oryza sativa L.		Oryza sativa (japonica cultivar-group); Oryza sativa subsp. japonica; Japanese rice; Japonica rice; Oryza sativa (japonica cultivar-group); Oryza sativa japonica; Oryza sativa ssp. japonica	
	Rank		Rank
species		no rank	
	Lineage		Lineage
cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliophyta; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; BOP clade; Oryzoideae; Oryzaceae; Oryzinae; Oryza		cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliophyta; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; BOP clade; Oryzoideae; Oryzaceae; Oryzinae; Oryza; Oryza sativa	
	Parent		Parent
Oryza () - (Rank: genus)		Oryza sativa (rice) - (Rank: species)	
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4527)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4530)	
4530	NCBI Taxonomy ID	39947	NCBI Taxonomy ID
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4530)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=39947)	
	is Taxon A an Intraspecies?		is Taxon B an Intraspecies?
Yes		Yes	
	Taxon A Description		Taxon B Description
Middle-late-flowering Tongil-type (japonica/indica hybrid) rice Milyang23 (116 days)		Early heading japonica rice H143 (78 days) and H75 (Hokkaido; Japan)	

GENOTYPIC CHANGE

	Generic Gene Name		UniProtKB Oryza sativa subsp. japonica
HD16		Q852L0 (http://www.uniprot.org/uniprot/Q852L0)	
	Synonyms		GenebankID or UniProtKB
CKI; EL1; Hd16; OsJ_12923; Os03g0793500; LOC_Os03g57940; OSJNBb0060J21.12		()	
	String		
39947.LOC_Os03g57940.1			
(http://string-db.org/newstring.cgi/show_network_section.pl?identifier=39947.LOC_Os03g57940.1)			
	Sequence Similarities		
Belongs to the protein kinase superfamily. CK1 Ser/Thr protein kinase family. Casein kinase I subfamily.			
	GO - Molecular Function		
GO:0005524 : ATP binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005524)			
GO:0004674 : protein serine/threonine kinase activity			
(https://www.ebi.ac.uk/QuickGO/term/GO:0004674)			

GO - Biological Process

GO:0009908 : flower development (<https://www.ebi.ac.uk/QuickGO/term/GO:0009908>)
 GO:0009740 : gibberellic acid mediated signaling pathway (<https://www.ebi.ac.uk/QuickGO/term/GO:0009740>)
 GO:0018105 : peptidyl-serine phosphorylation (<https://www.ebi.ac.uk/QuickGO/term/GO:0018105>)
 GO:0006897 : endocytosis (<https://www.ebi.ac.uk/QuickGO/term/GO:0006897>)
 GO:0048586 : regulation of long-day photoperiodism, flowering (<https://www.ebi.ac.uk/QuickGO/term/GO:0048586>)
 GO:0040008 : regulation of growth (<https://www.ebi.ac.uk/QuickGO/term/GO:0040008>)
 GO:0010476 : gibberellin mediated signaling pathway (<https://www.ebi.ac.uk/QuickGO/term/GO:0010476>)
 GO:0018107 : peptidyl-threonine phosphorylation (<https://www.ebi.ac.uk/QuickGO/term/GO:0018107>)

GO - Cellular Component

GO:0005737 : cytoplasm (<https://www.ebi.ac.uk/QuickGO/term/GO:0005737>)
 GO:0005634 : nucleus (<https://www.ebi.ac.uk/QuickGO/term/GO:0005634>)

Presumptive Null

No (<https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title>)

Molecular Type

Coding (<https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title>)

Aberration Type

SNP (<https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title>)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

G476C p.Gly159Ala in the serine/threonine kinase domain leading to non-functional protein

Experimental Evidence

Linkage Mapping (<https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Linkage Mapping^#gephebase-summary-title>)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	-	-	-

Main Reference

Natural variation in Early flowering1 contributes to early flowering in japonica rice under long days. (2014) (<https://pubmed.ncbi.nlm.nih.gov/23668360>)

Authors

Kwon CT; Yoo SC; Koo BH; Cho SH; Park JW; Zhang Z; Li J; Li Z; Paek NC

Abstract

Natural variation in heading-date genes enables rice, a short-day (SD) plant, to flower early under long-day (LD) conditions at high latitudes. Through analysis of heading-date quantitative trait loci (QTL) with F7 recombinant inbred lines from the cross of early heading 'H143' and late heading 'Milyang23 (M23)', we found a minor-effect Early Heading3 (EH3) QTL in the Hd16 region on chromosome 3. We found that Early flowering1 (EL1), encoding casein kinase I (CKI), is likely to be responsible for the EH3/Hd16 QTL, because a missense mutation occurred in the highly conserved serine/threonine kinase domain of EL1 in H143. A different missense mutation was found in the EL1 kinase domain in Koshihikari. In vitro kinase assays revealed that EL1/CKI in H143 and Koshihikari are non-functional. In F7:9 heterogeneous inbred family-near isogenic lines (HNILs), HNIL(H143) flowered 13 days earlier than HNIL(M23) in LD, but not in SD, in which EL1 mainly acts as a LD-dependent flowering repressor, down-regulating Ehd1 expression. In the world rice collection, two types of non-functional EL1 variants were found in japonica rice generally cultivated at high latitudes. These results indicate that natural variation in EL1 contributes to early heading for rice adaptation to LD in temperate and cooler regions.

© 2013 John Wiley & Sons Ltd.

Additional References

RELATED GEPHE

Related Genes

9 (DTH2, EARLY FLOWERING 3/Hd17, Hd1, Hd6a, PRR37 pseudoresponse regulator protein 37, se5, HEADING DATE 1, Ehd1 (Response regulator), Ghd7) (<https://www.gephebase.org/search-criteria?/or+Taxon ID=^4530^/and+Trait=Flowering time/or+Taxon ID=^39947^/and+Trait=Flowering time/and+groupHaplotypes=true#gephebase-summary-title>)

Related Haplotypes

1 ([https://www.gephebase.org/search-criteria?/or+Gene Gephebase=^Early flowering1 \(EL1\)^/and+Taxon ID=^4530^/or+Gene Gephebase=^Early flowering1 \(EL1\)^/and+Taxon ID=^39947^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/or+Gene Gephebase=^Early flowering1 (EL1)^/and+Taxon ID=^4530^/or+Gene Gephebase=^Early flowering1 (EL1)^/and+Taxon ID=^39947^#gephebase-summary-title))

EXTERNAL LINKS

COMMENTS

@GxE Non-functional variants are associated with cultivars of high latitudes

