

# GEPHE SUMMARY

	Gephebase Gene	GephelD
SCN8A (Nav1.6) ( <a href="https://www.gephebase.org/search-criteria?/and+Gene">https://www.gephebase.org/search-criteria?/and+Gene</a> Gephebase=^SCN8A (Nav1.6)^#gephebase-summary-title)	GP00001657	Main curator
	Entry Status	Prigent
Published		

## PHENOTYPIC CHANGE

	Trait Category		
Physiology ( <a href="https://www.gephebase.org/search-criteria?/and+Trait">https://www.gephebase.org/search-criteria?/and+Trait</a> Category="Physiology^#gephebase-summary-title")	Trait		
Xenobiotic resistance (TTX) ( <a href="https://www.gephebase.org/search-criteria?/and+Trait=Xenobiotic resistance (TTX)^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Trait=Xenobiotic resistance (TTX)^#gephebase-summary-title</a> )	Trait State in Taxon A		
TTX-Sensitive <i>Amphiesma pryeri</i>	Trait State in Taxon B		
TTX-Resistant natricine clade (Afronatrix; Rhabdophis; Xenocrophis)	Ancestral State		
Taxon A	Taxonomic Status		
Intergeneric or Higher ( <a href="https://www.gephebase.org/search-criteria?/and+Taxonomic">https://www.gephebase.org/search-criteria?/and+Taxonomic</a> Status="Intergeneric or Higher^#gephebase-summary-title")			
Taxon A	Latin Name	Taxon B	Latin Name
<i>Hebius pryeri</i> ( <a href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Hebius pryeri^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Hebius pryeri^#gephebase-summary-title</a> )	Natricinae ( <a href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Natricinae^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Natricinae^#gephebase-summary-title</a> )	-	Common Name
Pryer's keelback	Synonyms	Natricidae	Synonyms
<i>Amphiesma pryeri</i> ; <i>Tropidonotus pryeri</i> ; Pryer's keelback; <i>Amphiesma pryeri</i> (Boulenger, 1887)	Rank	subfamily	Rank
species	Lineage	cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Sarcopterygii; Dipnotetrapodomorpha; Tetrapoda; Amniota; Sauropsida; Sauria; Lepidosauria; Squamata; Bifurcata; Unidentata; Episquamata; Toxicofera; Serpentes; Colubroidea; Colubridae; Natricinae; <i>Hebius</i>	Lineage
<i>Hebius</i> () - (Rank: genus) ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 1591003">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 1591003</a> )	Parent	Colubridae (colubrid snakes) - (Rank: family) ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 8578">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 8578</a> )	NCBI Taxonomy ID
1159330 ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 1159330">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 1159330</a> )	NCBI Taxonomy ID	169862 ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 169862">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 169862</a> )	is Taxon B an Infraspecies?
		No	
No			

## GENOTYPIC CHANGE

SCN8A	Generic Gene Name	UniProtKB Afronatrix anoscopus A0A1BoZ7B0 ( <a href="http://www.uniprot.org/uniprot/A0A1BoZ7B0">http://www.uniprot.org/uniprot/A0A1BoZ7B0</a> )
-	Synonyms	GenebankID or UniProtKB
-	String	0
Belongs to the sodium channel (TC 1.A.1.10) family.	Sequence Similarities	
GO:0005244 : voltage-gated ion channel activity ( <a href="https://www.ebi.ac.uk/QuickGO/term/GO:0005244">https://www.ebi.ac.uk/QuickGO/term/GO:0005244</a> )	GO - Molecular Function	
GO:0005248 : voltage-gated sodium channel activity ( <a href="https://www.ebi.ac.uk/QuickGO/term/GO:0005248">https://www.ebi.ac.uk/QuickGO/term/GO:0005248</a> )		
GO:0034765 : regulation of ion transmembrane transport ( <a href="https://www.ebi.ac.uk/QuickGO/term/GO:0034765">https://www.ebi.ac.uk/QuickGO/term/GO:0034765</a> )	GO - Biological Process	

## GO - Cellular Component

GO:0001518 : voltage-gated sodium channel complex  
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0001518>)

Presumptive Null

No (<https://www.gephebase.org/search-criteria?/and+Presumptive+Null=%No%#gephebase-summary-title>)

Molecular Type

Coding (<https://www.gephebase.org/search-criteria?/and+Molecular+Type=%Coding%#gephebase-summary-title>)

Aberration Type

SNP (<https://www.gephebase.org/search-criteria?/and+Aberration+Type=%SNP%#gephebase-summary-title>)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

V1709I (reversion) in DIV (loss of resistance)

Experimental Evidence

Candidate Gene (<https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=%Candidate+Gene%#gephebase-summary-title>)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	-	-	-

Historical Contingency in a Multigene Family Facilitates Adaptive Evolution of Toxin Resistance. (2016) (<https://pubmed.ncbi.nlm.nih.gov/27291053>)

Main Reference

McGlothlin JW; Kobiela ME; Feldman CR; Castoe TA; Geffeney SL; Hanifin CT; Toledo G; Vonk FJ; Richardson MK; Brodie ED; Pfrender ME; Brodie ED

Authors

Novel adaptations must originate and function within an already established genome [1]. As a result, the ability of a species to adapt to new environmental challenges is predicted to be highly contingent on the evolutionary history of its lineage [2-6]. Despite a growing appreciation of the importance of historical contingency in the adaptive evolution of single proteins [7-11], we know surprisingly little about its role in shaping complex adaptations that require evolutionary change in multiple genes. One such adaptation, extreme resistance to tetrodotoxin (TTX), has arisen in several species of snakes through coevolutionary arms races with toxic amphibian prey, which select for TTX-resistant voltage-gated sodium channels (Nav) [12-16]. Here, we show that the relatively recent origins of extreme toxin resistance, which involve the skeletal muscle channel Nav1.4, were facilitated by ancient evolutionary changes in two other members of the same gene family. A substitution conferring TTX resistance to Nav1.7, a channel found in small peripheral neurons, arose in lizards ~4170 million years ago (mya) and was present in the common ancestor of all snakes. A second channel found in larger myelinated neurons, Nav1.6, subsequently evolved resistance in four different snake lineages beginning ~438 mya. Extreme TTX resistance has evolved at least five times within the past 12 million years via changes in Nav1.4, but only within lineages that previously evolved resistant Nav1.6 and Nav1.7. Our results show that adaptive protein evolution may be contingent upon enabling substitutions elsewhere in the genome, in this case, in paralogs of the same gene family.

Copyright © 2016 Elsevier Ltd. All rights reserved.

Additional References

## RELATED GEPHE

1 (SCN4A (Nav1.4)) (<https://www.gephebase.org/search-criteria?/or+Taxon+ID=%1159330%/and+Trait=Xenobiotic+resistance/or+Taxon+ID=%169862%/and+Trait=Xenobiotic+resistance/and+groupHaplotypes=true#gephebase-summary-title>)

Related Genes

Related Haplotypes

No matches found.

## EXTERNAL LINKS

## COMMENTS

Non-null mutation