

GEPHE SUMMARY

ATP4B (https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=^ATP4B^#gephebase-summary-title)	Gephebase Gene	GP00001929	GepheID
Published	Entry Status	Courtier	Main curator

PHENOTYPIC CHANGE

Physiology (https://www.gephebase.org/search-criteria?/and+Trait+Category=^Physiology^#gephebase-summary-title)	Trait Category
Digestion (absence of stomach) (https://www.gephebase.org/search-criteria?/and+Trait=^Digestion (absence of stomach)^#gephebase-summary-title)	Trait
presence of stomach and gastric acid production	Trait State in Taxon A
loss of stomach and no gastric acid production	Trait State in Taxon B
Taxon A	Ancestral State
Intergeneric or Higher (https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=^Intergeneric or Higher^#gephebase-summary-title)	Taxonomic Status

Taxon A #1	Latin Name
Oreochromis niloticus (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Oreochromis niloticus^#gephebase-summary-title)	
Nile tilapia	Common Name
Oreochromis nilotica; Tilapia nilotica; Nile tilapia; Oreochromis niloticus (Linnaeus, 1758)	Synonyms
species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Percomorpha; Ovalentaria; Cichlomorpha; Cichliformes; Cichlidae; African cichlids; Pseudocrenilabrinae; Oreochromini; Oreochromis	Lineage
Oreochromis () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 8139)	Parent
8128 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 8128)	NCBI Taxonomy ID
No	is Taxon A an Infrappecies?

Taxon B #1	Latin Name
Takifugu rubripes (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Takifugu rubripes^#gephebase-summary-title)	
torafugu	Common Name
Fugu rubripes; Sphaeroides rubripes; Tetraodon rubripes; torafugu; tiger puffer; Takifugu rubripes (Temminck & Schlegel, 1850)	Synonyms
species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Percomorpha; Eupercaria; Tetraodontiformes; Tetraodontoidei; Tetraodontoidea; Tetraodontidae; Takifugu	Lineage
Takifugu () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 31032)	Parent
31033 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 31033)	NCBI Taxonomy ID
No	is Taxon B an Infrappecies?

Taxon A #2	Latin Name
Gasterosteus aculeatus (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Gasterosteus aculeatus^#gephebase-summary-title)	
three-spined stickleback	Common Name
three-spined stickleback; three spined stickleback; Gasterosteus aculeatus Linnaeus, 1758	Synonyms
species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Percomorpha;	Lineage

Taxon B #2	Latin Name
Tetraodon nigroviridis (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Tetraodon nigroviridis^#gephebase-summary-title)	
spotted green pufferfish	Common Name
spotted green pufferfish; Tetraodon nigroviridis Marion de Proce, 1822	Synonyms
species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Percomorpha; Eupercaria; Tetraodontiformes; Tetraodontoidei; Tetraodontoidea; Tetraodontidae;	Lineage

Eupercaria; Perciformes; Cottioidei; Gasterosteales; Gasterosteidae; Gasterosteus
Parent
Gasterosteus () - (Rank: genus)
(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=69292>)
NCBI Taxonomy ID
69293
(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=69293>)
is Taxon A an Infrasppecies?
No

Tetraodon
Parent
Tetraodon () - (Rank: genus)
(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=47144>)
NCBI Taxonomy ID
99883
(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=99883>)
is Taxon B an Infrasppecies?
No

GENOTYPIC CHANGE

ATP4B	Generic Gene Name	P51164 (http://www.uniprot.org/uniprot/P51164)	UniProtKB Homo sapiens
ATP6B	Synonyms	0	GenebankID or UniProtKB
9606.ENSPO0000334216 (http://string-db.org/newstring.cgi/show_network_section.pl?identifier=9606.ENSPO0000334216)	String		
Belongs to the X(+)/potassium ATPases subunit beta family.	Sequence Similarities		
GO:0001671 : ATPase activator activity (https://www.ebi.ac.uk/QuickGO/term/GO:0001671)	GO - Molecular Function		
GO:0008900 : potassium:proton exchanging ATPase activity (https://www.ebi.ac.uk/QuickGO/term/GO:0008900)			
GO:0007155 : cell adhesion (https://www.ebi.ac.uk/QuickGO/term/GO:0007155)	GO - Biological Process		
GO:0034220 : ion transmembrane transport (https://www.ebi.ac.uk/QuickGO/term/GO:0034220)			
GO:0032496 : response to lipopolysaccharide (https://www.ebi.ac.uk/QuickGO/term/GO:0032496)			
GO:0030007 : cellular potassium ion homeostasis (https://www.ebi.ac.uk/QuickGO/term/GO:0030007)			
GO:0006883 : cellular sodium ion homeostasis (https://www.ebi.ac.uk/QuickGO/term/GO:0006883)			
GO:1990573 : potassium ion import across plasma membrane (https://www.ebi.ac.uk/QuickGO/term/GO:1990573)			
GO:0036376 : sodium ion export across plasma membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0036376)			
GO:0010243 : response to organonitrogen compound (https://www.ebi.ac.uk/QuickGO/term/GO:0010243)			
GO:0010248 : establishment or maintenance of transmembrane electrochemical gradient (https://www.ebi.ac.uk/QuickGO/term/GO:0010248)			
GO:0005886 : plasma membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0005886)	GO - Cellular Component		
GO:0005890 : sodium:potassium-exchanging ATPase complex (https://www.ebi.ac.uk/QuickGO/term/GO:0005890)			
Yes (https://www.gephebase.org/search-criteria?/and+Presumptive Null=~Yes^#gephebase-summary-title)			Presumptive Null
Gene Loss (https://www.gephebase.org/search-criteria?/and+Molecular Type=~Gene Loss^#gephebase-summary-title)			Molecular Type
Deletion (https://www.gephebase.org/search-criteria?/and+Aberration Type=~Deletion^#gephebase-summary-title)			Aberration Type
-			Deletion Size
Absence of the gene in the genome sequence - high synteny			Molecular Details of the Mutation
Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=~Candidate Gene^#gephebase-summary-title)			Experimental Evidence
Recurrent gene loss correlates with the evolution of stomach phenotypes in gnathostome history. (2014) (https://pubmed.ncbi.nlm.nih.gov/24307675)			Main Reference
Castro LF; GonÁsalves O; Mazan S; Tay BH; Venkatesh B; Wilson JM			Authors
The stomach, a hallmark of gnathostome evolution, represents a unique anatomical innovation characterized by the presence of acid- and pepsin-secreting glands. However, the occurrence of these glands in gnathostome species is not universal; in the nineteenth century the French zoologist Cuvier first noted that some teleosts lacked a stomach. Strikingly, Holocephali (chimaeras), dipnoids (lungfish) and monotremes (egg-laying mammals) also lack acid secretion and a gastric cellular phenotype. Here, we test the hypothesis that loss of the gastric phenotype is correlated with the loss of key gastric genes. We investigated species from all the main gnathostome lineages and show the specific contribution of gene loss to the widespread distribution of the agastric condition. We establish that the stomach loss correlates with the persistent and complete absence of the gastric function gene kit--H(+)/K(+)-ATPase (Atp4A and Atp4B) and pepsinogens (Pga, Pgc, Cym)--in the analysed species. We also find that in gastric species the pepsinogen gene complement varies significantly (e.g. two to four in teleosts and tens in some mammals) with multiple events of pseudogenization identified in various lineages. We propose that relaxation of purifying selection in pepsinogen genes and possibly proton pump genes in response to dietary changes led to the numerous independent events of stomach loss in gnathostome history. Significantly, the absence of the gastric genes predicts that reinvention of the stomach in agastric lineages would be highly improbable, in line with Dollo's principle.			Abstract
			Additional References

RELATED GEPHE

Related Genes

4 (ATP4A, pepsinogen A1, pepsinogen A2, pepsinogen A3) (<https://www.gephebase.org/search-criteria?/or+Taxon ID=^8128^/and+Trait=Digestion/or+Taxon ID=^69293^/and+Trait=Digestion/or+Taxon ID=^31033^/and+Trait=Digestion/or+Taxon ID=^99883^/and+Trait=Digestion/and+groupHaplotypes=true#gephebase-summary-title>)

Related Haplotypes

2 (<https://www.gephebase.org/search-criteria?/or+Gene Gephabase=^ATP4B^/and+Taxon ID=^8128^/or+Gene Gephabase=^ATP4B^/and+Taxon ID=^69293^/or+Gene Gephabase=^ATP4B^/and+Taxon ID=^31033^/or+Gene Gephabase=^ATP4B^/and+Taxon ID=^99883^#gephebase-summary-title>)

EXTERNAL LINKS

COMMENTS

Not clear if this is independent evolution in *Tetraodon nigroviridis* and in *Takifugu rubripes* (no detailed phylogenetic tree in the paper)