

GEPHE SUMMARY

<p>Fads2 (<a +fads2+"#gephebase-summary-title"="" href="https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=">https://www.gephebase.org/search-criteria?/and+Gene+Gephebase="+Fads2+"#gephebase-summary-title)</p> <p>Published</p>	<p>Gephebase Gene</p> <p>Entry Status</p>	<p>GP00002053</p> <p>Courtier</p>	<p>GepheID</p> <p>Main curator</p>
---	---	-----------------------------------	------------------------------------

PHENOTYPIC CHANGE

<p>Physiology (<a +physiology+"#gephebase-summary-title"="" href="https://www.gephebase.org/search-criteria?/and+Trait+Category=">https://www.gephebase.org/search-criteria?/and+Trait+Category="+Physiology+"#gephebase-summary-title)</p> <p>Fatty acid metabolism (fatty acid desaturation) (<a (fatty="" +fatty="" acid="" desaturation)+"#gephebase-summary-title"="" href="https://www.gephebase.org/search-criteria?/and+Trait=" metabolism="">https://www.gephebase.org/search-criteria?/and+Trait="+Fatty acid metabolism (fatty acid desaturation)+"#gephebase-summary-title)</p> <p>Low physiological ability to survive in freshwater DHA free diets</p> <p>High physiological ability to survive in freshwater DHA free diets</p> <p>Taxon A</p> <p>Interspecific (<a +interspecific+"#gephebase-summary-title"="" href="https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=">https://www.gephebase.org/search-criteria?/and+Taxonomic+Status="+Interspecific+"#gephebase-summary-title)</p>	<p>Trait Category</p> <p>Trait</p> <p>Trait State in Taxon A</p> <p>Trait State in Taxon B</p> <p>Ancestral State</p> <p>Taxonomic Status</p>
--	---

<p>Pungitius pungitius (<a +pungitius="" href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=" pungitius+"#gephebase-summary-title"="">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="+Pungitius pungitius+"#gephebase-summary-title)</p> <p>ninespine stickleback</p> <p>Gasterosteus pungitius; ninespine stickleback; Gasterosteus pungitius Linnaeus, 1758; Pungitius pungitius (Linnaeus, 1758)</p> <p>species</p> <p>cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Perciformes; Cottioidei; Gasterosteales; Gasterosteidae; Pungitius</p> <p>Pungitius () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=134919)</p> <p>134920 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=134920)</p> <p>No is Taxon A an Intraspecies?</p>	<p>Taxon A</p> <p>Latin Name</p> <p>Common Name</p> <p>Synonyms</p> <p>Rank</p> <p>Lineage</p> <p>Parent</p> <p>NCBI Taxonomy ID</p>
---	--

<p>Pungitius tymensis (<a +pungitius="" href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=" tymensis+"#gephebase-summary-title"="">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="+Pungitius tymensis+"#gephebase-summary-title)</p> <p>Sakhalin stickleback</p> <p>Gasterosteus tymensis; Sakhalin stickleback; Gasterosteus tymensis Nikolskii, 1889; Pungitius tymensis (Nikolskii, 1889)</p> <p>species</p> <p>cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Perciformes; Cottioidei; Gasterosteales; Gasterosteidae; Pungitius</p> <p>Pungitius () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=1480064)</p> <p>1480064 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=1480064)</p> <p>No is Taxon B an Intraspecies?</p>	<p>Taxon B #1</p> <p>Latin Name</p> <p>Common Name</p> <p>Synonyms</p> <p>Rank</p> <p>Lineage</p> <p>Parent</p> <p>NCBI Taxonomy ID</p>
--	---

<p>Pungitius kaibarae (<a +pungitius="" href="https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=" kaibarae+"#gephebase-summary-title"="">https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms="+Pungitius kaibarae+"#gephebase-summary-title)</p> <p>-</p> <p>Amur stickleback; Pygosteus kaibarae; Pungitius kaibarae (Tanaka, 1915); Pygosteus kaibarae Tanaka, 1915</p> <p>species</p> <p>cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Deuterostomia; Chordata; Craniata; Vertebrata; Gnathostomata; Teleostomi; Euteleostomi; Actinopterygii; Actinopteri; Neopterygii; Teleostei; Osteoglossocephalai; Clupeocephala; Euteleostomorpha; Neoteleostei; Eurypterygia; Ctenosquamata; Acanthomorpha; Euacanthomorpha; Perciformes; Cottioidei; Gasterosteales; Gasterosteidae; Pungitius</p>	<p>Taxon B #2</p> <p>Latin Name</p> <p>Common Name</p> <p>Synonyms</p> <p>Rank</p> <p>Lineage</p>
--	---

Eupercaria; Perciformes; Cottioidei; Gasterosteales; Gasterosteidae; Pungitius

Parent

Pungitius () - (Rank: genus)

(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=134919>)

NCBI Taxonomy ID

300318

(<https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=300318>)

is Taxon B an Intraspecies?

No

GENOTYPIC CHANGE

fads2	Generic Gene Name	Q9DEX7 (http://www.uniprot.org/uniprot/Q9DEX7)	UniProtKB Danio rerio
Fadsd6; DRD5/D6; wu:fb64c04; wu:fb69e08; zgc:112502; fadsd6	Synonyms	()	GenebankID or UniProtKB
7955.ENSADARP0000022396 (http://string-db.org/newstring.cgi/show_network_section.pl?identifier=7955.ENSADARP0000022396)	String		
Belongs to the fatty acid desaturase type 1 family.	Sequence Similarities		
GO:0016213 : linoleoyl-CoA desaturase activity (https://www.ebi.ac.uk/QuickGO/term/GO:0016213)	GO - Molecular Function		
GO:0006636 : unsaturated fatty acid biosynthetic process (https://www.ebi.ac.uk/QuickGO/term/GO:0006636)	GO - Biological Process		
GO:0001889 : liver development (https://www.ebi.ac.uk/QuickGO/term/GO:0001889)	GO - Cellular Component		
GO:0016021 : integral component of membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0016021)			
GO:0005789 : endoplasmic reticulum membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0005789)			
No (#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Presumptive Null="No">#gephebase-summary-title)			Presumptive Null
Gene Amplification (<a amplification"="" gene="" href="https://www.gephebase.org/search-criteria?/and+Molecular Type=">#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Molecular Type="Gene Amplification">#gephebase-summary-title)			Molecular Type
Insertion (#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Aberration Type="Insertion">#gephebase-summary-title)			Aberration Type
1-10 kb			Insertion Size
higher number of copies of Fads2.			Molecular Details of the Mutation
Candidate Gene (#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Experimental Evidence="Candidate Gene">#gephebase-summary-title)			Experimental Evidence
A key metabolic gene for recurrent freshwater colonization and radiation in fishes. (2019) (https://pubmed.ncbi.nlm.nih.gov/31147520)			Main Reference
Ishikawa A; Kabeya N; Ikeya K; Kakioka R; Cech JN; Osada N; Leal MC; Inoue J; Kume M; Toyoda A; Tezuka A; Nagano AJ; Yamasaki YY; Suzuki Y; Kokita T; Takahashi H; Lucek K; Marques D; Takehana Y; Naruse K; Mori S; Monroig O; Ladd N; Schubert CJ; Matthews B; Peichel CL; Seehausen O; Yoshizaki G; Kitano J			Authors
Colonization of new ecological niches has triggered large adaptive radiations. Although some lineages have made use of such opportunities, not all do so. The factors causing this variation among lineages are largely unknown. Here, we show that deficiency in docosahexaenoic acid (DHA), an essential 18:3 fatty acid, can constrain freshwater colonization by marine fishes. Our genomic analyses revealed multiple independent duplications of the fatty acid desaturase gene Fads2 in stickleback lineages that subsequently colonized and radiated in freshwater habitats, but not in close relatives that failed to colonize. Transgenic manipulation of Fads2 in marine stickleback increased their ability to synthesize DHA and survive on DHA-deficient diets. Multiple freshwater ray-finned fishes also show a convergent increase in Fads2 copies, indicating its key role in freshwater colonization.			Abstract

Copyright © 2019 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works.

Additional References

RELATED GEPHE

No matches found.

Related Genes

No matches found.

Related Haplotypes

EXTERNAL LINKS

COMMENTS

Both *P. tymensis* and *P. kaibarae* are entirely freshwater species. *Pungitius sinensis*, which inhabits both freshwater and brackish environments; has *Fads2* copy numbers intermediate between those of *P. pungitius* and freshwater nine-spined sticklebacks