

GEPHE SUMMARY

ABCA2 (https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=^ABCA2^#gephebase-summary-title)	Gephebase Gene	GP00002054	GepheID
Published	Entry Status	Courtier	Main curator

PHENOTYPIC CHANGE

Physiology (https://www.gephebase.org/search-criteria?/and+Trait+Category=^Physiology^#gephebase-summary-title)	Trait Category		
Xenobiotic resistance (insecticide; Bt Cry2Ab toxin) (https://www.gephebase.org/search-criteria?/and+Trait=^Xenobiotic+resistance+(insecticide;+Bt+Cry2Ab+toxin)^#gephebase-summary-title)	Trait		
Trichoplusia ni - Bt-Cry2Ab susceptible	Trait State in Taxon A		
Trichoplusia ni - Bt-Cry2Ab resistant	Trait State in Taxon B		
Taxon A	Ancestral State		
Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=^Intraspecific^#gephebase-summary-title)	Taxonomic Status		
		Taxon A	Taxon B
	Latin Name		Latin Name
Trichoplusia ni (https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=^Trichoplusia+ni^#gephebase-summary-title)		Trichoplusia ni (https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=^Trichoplusia+ni^#gephebase-summary-title)	
	Common Name		Common Name
cabbage looper		cabbage looper	
	Synonyms		Synonyms
cabbage looper; Trichoplusia ni (Hubner, 1803); Trichoplusia ni; Trichopulsia ni		cabbage looper; Trichoplusia ni (Hubner, 1803); Trichoplusia ni; Trichopulsia ni	
	Rank		Rank
species		species	
	Lineage		Lineage
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphimesenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Dityrsia; Obtectomera; Noctuoidea; Noctuidae; Plusiinae; Trichoplusia		cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphimesenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Dityrsia; Obtectomera; Noctuoidea; Noctuidae; Plusiinae; Trichoplusia	
	Parent		Parent
Trichoplusia () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=7110)		Trichoplusia () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=7110)	
	NCBI Taxonomy ID		NCBI Taxonomy ID
7111 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=7111)		7111 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=7111)	
	is Taxon A an Intraspecies?		is Taxon B an Intraspecies?
No		No	

GENOTYPIC CHANGE

ABCA2	Generic Gene Name	UniProtKB Helicoverpa armigera A0A0S0G7V0 (http://www.uniprot.org/uniprot/A0A0S0G7V0)
-	Synonyms	GenebankID or UniProtKB 0
-	String	
-	Sequence Similarities	
-	GO - Molecular Function	
GO:0005524 : ATP binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005524)		
GO:0042626 : ATPase activity, coupled to transmembrane movement of substances (https://www.ebi.ac.uk/QuickGO/term/GO:0042626)		
-	GO - Biological Process	
-	GO - Cellular Component	
GO:0016021 : integral component of membrane		

(<https://www.ebi.ac.uk/QuickGO/term/GO:0016021>)

Presumptive Null

Yes ([https://www.gephebase.org/search-criteria?/and+Presumptive Null=^Yes^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Presumptive+Null=^Yes^#gephebase-summary-title))

Molecular Type

Coding ([https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Molecular+Type=^Coding^#gephebase-summary-title))

Aberration Type

Insertion ([https://www.gephebase.org/search-criteria?/and+Aberration Type=^Insertion^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Aberration+Type=^Insertion^#gephebase-summary-title))

Insertion Size

1-10 kb

Molecular Details of the Mutation

insertion of a transposon Tntransib (2581 bp) in ABCA2 which changes splicing sites in the transcript and lead to an indel in the coding region: change of 1551VETLAHALGFLRHLDKRR1567 into 1551AHWGK- LYGSNTQN1563

Experimental Evidence

Linkage Mapping ([https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Linkage Mapping^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=^Linkage+Mapping^#gephebase-summary-title))

Main Reference

Mutation of ABC transporter ABCA2 confers resistance to Bt toxin Cry2Ab in *Trichoplusia ni*. (2019) (<https://pubmed.ncbi.nlm.nih.gov/31422154>)

Authors

Yang X; Chen W; Song X; Ma X; Cotto-Rivera RO; Kain W; Chu H; Chen YR; Fei Z; Wang P

Abstract

Insecticidal proteins from *Bacillus thuringiensis* (Bt) are the primary recombinant proteins expressed in transgenic crops (Bt-crops) to confer insect resistance. Development of resistance to Bt toxins in insect populations threatens the sustainable application of Bt-crops in agriculture. The Bt toxin Cry2Ab is a major insecticidal protein used in current Bt-crops, and resistance to Cry2Ab has been selected in several insects, including the cabbage looper, *Trichoplusia ni*. In this study, the Cry2Ab resistance gene in *T. ni* was mapped to Chromosome 17 by genetic linkage analyses using a whole genome resequencing approach, and was then finely mapped using RNA-seq-based bulked segregant analysis (BSA) and amplicon sequencing (AmpSeq)-based fine linkage mapping to a locus containing two genes, ABCA1 and ABCA2. Mutations in ABCA1 and ABCA2 in Cry2Ab resistant *T. ni* were identified by both genomic DNA and cDNA sequencing. Analysis of the expression of ABCA1 and ABCA2 in *T. ni* larvae indicated that ABCA2 is abundantly expressed in the larval midgut, but ABCA1 is not a midgut-expressed gene. The mutation in ABCA2 in Cry2Ab resistant *T. ni* was identified to be an insertion of a transposon Tntransib in ABCA2. For confirmation of ABCA2 as the Cry2Ab-resistance gene, *T. ni* mutants with frameshift mutations in ABCA1 and ABCA2 were generated by CRISPR/Cas9 mutagenesis. Bioassays of the *T. ni* mutants with Cry2Ab verified that the mutations of ABCA1 did not change larval susceptibility to Cry2Ab, but the ABCA2 mutants were highly resistant to Cry2Ab. Genetic complementation test of the ABCA2 allele in Cry2Ab resistant *T. ni* with an ABCA2 mutant generated by CRISPR/Cas9 confirmed that the ABCA2 mutation in the Cry2Ab resistant strain confers the resistance. The results from this study confirmed that ABCA2 is essential for the toxicity of Cry2Ab in *T. ni* and mutation of ABCA2 confers the resistance to Cry2Ab in the resistant *T. ni* strain derived from a Bt resistant greenhouse population.

Copyright © 2019 Elsevier Ltd. All rights reserved.

Additional References

RELATED GEPHE

Related Genes

1 (ABCC2) ([https://www.gephebase.org/search-criteria?/or+Taxon ID=^7111^/and+Trait=Xenobiotic resistance/and+groupHaplotypes=true#gephebase-summary-title](https://www.gephebase.org/search-criteria?/or+Taxon+ID=^7111^/and+Trait=Xenobiotic+resistance/and+groupHaplotypes=true#gephebase-summary-title))

Related Haplotypes

No matches found.

EXTERNAL LINKS

COMMENTS

@TE @Splicing