

GEPHE SUMMARY

	Gephebase Gene		GepheID
flavonoid 3'-hydroxylase (F3'H) (https://www.gephebase.org/search-criteria?/and+Gene)		GP00002087	
Gephebase="flavonoid 3'-hydroxylase (F3'H)"#gephebase-summary-title)			Main curator
Published	Entry Status	Courtier	

PHENOTYPIC CHANGE

	Trait Category	
Morphology (https://www.gephebase.org/search-criteria?/and+Trait)		
Category="Morphology"#gephebase-summary-title)	Trait	
Coloration (flowers) (<a "="" href="https://www.gephebase.org/search-criteria?/and+Trait=">https://www.gephebase.org/search-criteria?/and+Trait=")		
(flowers)"#gephebase-summary-title)	Trait State in Taxon A	
Ipomoea nil - bright blue flowers		
	Trait State in Taxon B	
Ipomoea nil - magenta mutant - stable reddish flowers		
	Ancestral State	
Taxon A		
	Taxonomic Status	
Domesticated (https://www.gephebase.org/search-criteria?/and+Taxonomic)		
Status="Domesticated"#gephebase-summary-title)		

Taxon A	Latin Name	Taxon B	Latin Name
Ipomoea nil (https://www.gephebase.org/search-criteria?/and+Taxon)		Ipomoea nil (https://www.gephebase.org/search-criteria?/and+Taxon)	
nil"#gephebase-summary-title)	Common Name	nil"#gephebase-summary-title)	Common Name
Japanese morning glory		Japanese morning glory	
	Synonyms		Synonyms
Convolvulus nil; Pharbitis nil; Japanese morning glory; qian niu; Convolvulus nil L., 1762;		Convolvulus nil; Pharbitis nil; Japanese morning glory; qian niu; Convolvulus nil L., 1762;	
Ipomoea nil (L.) Roth, 1797; Pharbitis nil (L.) Choisy, 1834		Ipomoea nil (L.) Roth, 1797; Pharbitis nil (L.) Choisy, 1834	
	Rank		Rank
species		species	
	Lineage		Lineage
cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta;		cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta;	
Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliopsida; Mesangiospermae;		Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliopsida; Mesangiospermae;	
eudicotyledons; Gunneridae; Pentapetalae; asterids; lamiids; Solanales; Convolvulaceae;		eudicotyledons; Gunneridae; Pentapetalae; asterids; lamiids; Solanales; Convolvulaceae;	
Ipomoeae; Ipomoea		Ipomoeae; Ipomoea	
	Parent		Parent
Ipomoea () - (Rank: genus)		Ipomoea () - (Rank: genus)	
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4119)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4119)	
	NCBI Taxonomy ID		NCBI Taxonomy ID
35883		35883	
(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=35883)		(https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=35883)	
	is Taxon A an Intraspecies?		is Taxon B an Intraspecies?
No		No	

GENOTYPIC CHANGE

	Generic Gene Name	UniProtKB Arabidopsis thaliana
CYP75B1		Q9SD85 (http://www.uniprot.org/uniprot/Q9SD85)
	Synonyms	GenebankID or UniProtKB
CYP75B1; CYTOCHROME P450 75B1; D501; F13G24.190; F13G24_190; F3'H;		
FLAVONOID 3'-HYDROXYLASE; TRANSPARENT TESTA 7; TT7; At5g07990		
	String	
3702.AT5G07990.1		
(http://string-db.org/newstring.cgi/show_network_section.pl?identifier=3702.AT5G07990.1)		
	Sequence Similarities	
Belongs to the cytochrome P450 family.		
	GO - Molecular Function	
GO:0020037 : heme binding (https://www.ebi.ac.uk/QuickGO/term/GO:0020037)		
GO:0005506 : iron ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005506)		
GO:0016709 : oxidoreductase activity, acting on paired donors, with incorporation or		
reduction of molecular oxygen, NAD(P)H as one donor, and incorporation of one atom of		
oxygen (https://www.ebi.ac.uk/QuickGO/term/GO:0016709)		

GO - Biological Process

GO:0009733 : response to auxin (<https://www.ebi.ac.uk/QuickGO/term/GO:0009733>)

GO:0009813 : flavonoid biosynthetic process
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0009813>)

GO - Cellular Component

GO:0016021 : integral component of membrane
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0016021>)

GO:0016020 : membrane (<https://www.ebi.ac.uk/QuickGO/term/GO:0016020>)

GO:0005789 : endoplasmic reticulum membrane
 (<https://www.ebi.ac.uk/QuickGO/term/GO:0005789>)

Presumptive Null

Yes ([https://www.gephebase.org/search-criteria?/and+Presumptive Null=~Yes^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Presumptive+Null=~Yes^#gephebase-summary-title))

Molecular Type

Coding ([https://www.gephebase.org/search-criteria?/and+Molecular Type=~Coding^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Molecular+Type=~Coding^#gephebase-summary-title))

Aberration Type

SNP ([https://www.gephebase.org/search-criteria?/and+Aberration Type=~SNP^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Aberration+Type=~SNP^#gephebase-summary-title))

SNP Coding Change

Nonsense

Molecular Details of the Mutation

nonsense mutation caused by a single C to T base transition generating the stop codon TGA.

Experimental Evidence

Candidate Gene ([https://www.gephebase.org/search-criteria?/and+Experimental Evidence=~Candidate Gene^#gephebase-summary-title](https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=~Candidate+Gene^#gephebase-summary-title))

	Taxon A	Taxon B	Position
Codon	CGA	TGA	-
Amino-acid	Arg	STP	-

Main Reference

Spontaneous mutations of the flavonoid 3'-hydroxylase gene conferring reddish flowers in the three morning glory species. (2003) (<https://pubmed.ncbi.nlm.nih.gov/14581624>)

Authors

Hoshino A; Morita Y; Choi JD; Saito N; Toki K; Tanaka Y; Iida S

Abstract

Among the Ipomoea plants, both Ipomoea nil and Ipomoea tricolor display bright blue flowers, and Ipomoea purpurea exhibits dark purple flowers. While all of these flowers contain cyanidin-based anthocyanin pigments, the mutants of I. nil, I. purpurea, and I. tricolor carrying the magenta, pink, and fuchsia alleles, respectively, produce reddish flowers containing pelargonidin derivatives, and all of them are deficient in the gene for flavonoid 3'-hydroxylase (F3'H). The magenta allele in I. nil is a nonsense mutation caused by a single C to T base transition generating the stop codon TGA, and the cultivar Violet carries the same mutation. Several tested pink mutants in I. purpurea carry inserts of the 0.55-kb DNA transposable element Tip201 belonging to the Ac/Ds superfamily at the identical site. No excision of Tip201 from the F3'H gene could be detected, and both splicing and polyadenylation patterns of the F3'H transcripts were affected by the Tip201 integration. The fuchsia allele in I. tricolor is a single T insertion generating the stop codon TAG, and the accumulation of the F3'H transcripts was drastically reduced by the nonsense-mediated RNA decay. Spontaneous mutations in Ipomoea, including a possible founder mutation in the pink allele, are also discussed.

Additional References

RELATED GEPHE

Related Genes

2 (MYB1, WDR1) ([https://www.gephebase.org/search-criteria?/or+Taxon ID=~35883^/and+Trait=Coloration/and+groupHaplotypes=true#gephebase-summary-title](https://www.gephebase.org/search-criteria?/or+Taxon+ID=~35883^/and+Trait=Coloration/and+groupHaplotypes=true#gephebase-summary-title))

Related Haplotypes

No matches found.

EXTERNAL LINKS

COMMENTS