

GEPHE SUMMARY

para (kdr) (https://www.gephebase.org/search-criteria?/and+Gene Gephebase='para (kdr)'#gephebase-summary-title)	Gephebase Gene	GP00002505	GepheID
	Entry Status	Courtier	Main curator
Published			

PHENOTYPIC CHANGE

Trait Category			
Physiology (https://www.gephebase.org/search-criteria?/and+Trait Category='Physiology'#gephebase-summary-title)	Trait		
Xenobiotic resistance (insecticide) (https://www.gephebase.org/search-criteria?/and+Trait='Xenobiotic resistance (insecticide)'#gephebase-summary-title)	Trait State in Taxon A		
Sitophilus zeamais	Trait State in Taxon B		
Sitophilus zeamais - resistant	Ancestral State		
Taxon A	Taxonomic Status		
Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic Status='Intraspecific'#gephebase-summary-title)			
Taxon A	Latin Name	Taxon B	Latin Name
Sitophilus zeamais (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms='Sitophilus zeamais'#gephebase-summary-title)		Sitophilus zeamais (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms='Sitophilus zeamais'#gephebase-summary-title)	
maize weevil	Common Name	maize weevil	Common Name
Sitophilus oryzae zeamais; maize weevil; Sitophilus zeamais Motschulsky, 1855	Synonyms	Sitophilus oryzae zeamais; maize weevil; Sitophilus zeamais Motschulsky, 1855	Synonyms
species	Rank	species	Rank
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Coleoptera; Polyphaga; Cucujiformia; Curculionoidea; Curculionidae; Dryophthorinae; Sitophilus	Lineage	cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Coleoptera; Polyphaga; Cucujiformia; Curculionoidea; Curculionidae; Dryophthorinae; Sitophilus	Lineage
Sitophilus () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 7045)	Parent	Sitophilus () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 7045)	Parent
7047 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 7047)	NCBI Taxonomy ID	7047 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 7047)	NCBI Taxonomy ID
No	is Taxon A an Infraspecies?	No	is Taxon B an Infraspecies?

GENOTYPIC CHANGE

Generic Gene Name			
para	Synonyms	P35500 (http://www.uniprot.org/uniprot/P35500)	UniProtKB Drosophila melanogaster
bas; bss; CG9907; Dmel\CG9907; DmNav; DmNav1; DmNa[[v]]; DmNa[[V]]; DmNa[[v]]1; l(1)14Da; l(1)ESHS48; lincRNA.S9469; Nav1; Ocd; olfD; par; sbl; sbl-1; Shu; Shudderer	String	0	GenebankID or UniProtKB
7227.FBpp0303597 (http://string-db.org/newstring_cgi/show_network_section.pl?identifier= 7227.FBpp0303597)	Sequence Similarities		
Belongs to the sodium channel (TC 1.A.1.10) family. Para subfamily.	GO - Molecular Function		
GO:0005509 : calcium ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005509)			
GO:0005244 : voltage-gated ion channel activity (https://www.ebi.ac.uk/QuickGO/term/GO:0005244)			
GO:0005248 : voltage-gated sodium channel activity (https://www.ebi.ac.uk/QuickGO/term/GO:0005248)			
GO:0005272 : sodium channel activity			

(<https://www.ebi.ac.uk/QuickGO/term/GO:0005272>)

GO - Biological Process

GO:0045433 : male courtship behavior, veined wing generated song production

(<https://www.ebi.ac.uk/QuickGO/term/GO:0045433>)

GO:0001666 : response to hypoxia (<https://www.ebi.ac.uk/QuickGO/term/GO:0001666>)

GO:0009612 : response to mechanical stimulus

(<https://www.ebi.ac.uk/QuickGO/term/GO:0009612>)

GO:0034765 : regulation of ion transmembrane transport

(<https://www.ebi.ac.uk/QuickGO/term/GO:0034765>)

GO:0035725 : sodium ion transmembrane transport

(<https://www.ebi.ac.uk/QuickGO/term/GO:0035725>)

GO:0007638 : mechanosensory behavior

(<https://www.ebi.ac.uk/QuickGO/term/GO:0007638>)

GO:0060078 : regulation of postsynaptic membrane potential

(<https://www.ebi.ac.uk/QuickGO/term/GO:0060078>)

GO - Cellular Component

GO:0005887 : integral component of plasma membrane

(<https://www.ebi.ac.uk/QuickGO/term/GO:0005887>)

GO:0001518 : voltage-gated sodium channel complex

(<https://www.ebi.ac.uk/QuickGO/term/GO:0001518>)

Presumptive Null

No (<https://www.gephebase.org/search-criteria?/and+Presumptive+Null=%No%#gephebase-summary-title>)

Molecular Type

Coding (<https://www.gephebase.org/search-criteria?/and+Molecular+Type=%Coding%#gephebase-summary-title>)

Aberration Type

SNP (<https://www.gephebase.org/search-criteria?/and+Aberration+Type=%SNP%#gephebase-summary-title>)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

T929I

Experimental Evidence

Candidate Gene (<https://www.gephebase.org/search-criteria?/and+Experimental+Evidence=%Candidate+Gene%#gephebase-summary-title>)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Thr	Ile	929

Main Reference

Pyrethroid resistance in *Sitophilus zeamais* is associated with a mutation (T929I) in the voltage-gated sodium channel. (2011) (<https://pubmed.ncbi.nlm.nih.gov/21496128>)

Authors

Araújo RA; Williamson MS; Bass C; Field LM; Duce IR

Abstract

The maize weevil, *Sitophilus zeamais*, is the most important pest affecting stored grain in Brazil and its control relies heavily on the use of insecticides. The intensive use of compounds such as the pyrethroids has led to the emergence of resistance, and previous studies have suggested that resistance to both pyrethroids and 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane (DDT) may result from reduced sensitivity of the insecticide target, the voltage-gated sodium channel. To identify the molecular mechanisms underlying pyrethroid resistance in *S. zeamais*, the domain II region of the voltage-gated sodium channel (para-orthologue) gene was amplified by PCR and sequenced from susceptible and resistant laboratory *S. zeamais* strains that were selected with a discriminating dose of DDT. A single point mutation, T929I, was found in the para gene of the resistant *S. zeamais* populations and its presence in individual weevils was strongly associated with survival after DDT exposure. This is the first identification of a target-site resistance mutation in *S. zeamais* and unusually it is a super-kdr type mutation occurring in the absence of the more common kdr (L1014F) substitution. A high-throughput assay based on TaqMan single nucleotide polymorphism genotyping was developed for sensitive detection of the mutation and used to screen field-collected strains of *S. zeamais*. This showed that the mutation is present at low frequency in field populations and is a useful tool for informing control strategies.

© 2011 The Authors. Insect Molecular Biology © 2011 The Royal Entomological Society.

Additional References

Molecular biology of insect sodium channels and pyrethroid resistance. (2014) (<https://pubmed.ncbi.nlm.nih.gov/24704279>)

RELATED GEPHE

Related Genes

No matches found.

Related Haplotypes

No matches found.

EXTERNAL LINKS

COMMENTS

It is a super-kdr type mutation occurring in the absence of the more common L1014F substitution.