GEPHE SUMMARY

Gephebase Gene
para (kdr) (https://www.gephebase.org/search-criteria?/and+Gene Gephebase=^para (kdr)^#gephebase-summary-title)

GepheID
GP00002630

Entry Status
Published

Main curator
Courtier

PHENOTYPIC CHANGE

Trait Category
Physiology (https://www.gephebase.org/search-criteria?/and+Trait Category=^Physiology^#gephebase-summary-title)

Trait
Xenobiotic resistance (insecticide) (https://www.gephebase.org/search-criteria?/and+Trait=^Xenobiotic resistance (insecticide)^#gephebase-summary-title)

Trait State in Taxon A
Plutella xylostella - sensitive

Trait State in Taxon B
Plutella xylostella - resistant in China

Ancestral State
Taxon A

Taxonomic Status
Intraspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic Status=^Intraspecific^#gephebase-summary-title)

Taxon A

Latin Name
Plutella xylostella (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Plutella xylostella^#gephebase-summary-title)

Common Name
diamondback moth

Synonyms
diamondback moth; cabbage moth; Plutella xylostella (Linnaeus, 1758); Putella xylostella

Rank
species

Lineage
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Yponomeutoidea; Plutellidae; Plutella

Parent
Plutella () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 51654)

NCBI Taxonomy ID
51655 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 51655)

is Taxon A an Infraspecies?
No

Taxon B

Latin Name
Plutella xylostella (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Plutella xylostella^#gephebase-summary-title)

Common Name
diamondback moth

Synonyms
diamondback moth; cabbage moth; Plutella xylostella (Linnaeus, 1758); Putella xylostella

Rank
species

Lineage
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Yponomeutoidea; Plutellidae; Plutella

Parent
Plutella () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 51654)

NCBI Taxonomy ID
51655 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 51655)

is Taxon B an Infraspecies?
No

GENOTYPIC CHANGE

Generic Gene Name
para

Synonyms
bas; bss; CG9907; Dmel\CG9907; DmNav; DmNav1; DmNa[[v]]; DmNa[[V]]; DmNa[[v]]1; l(1)14Da; l(1)ESHS48; lincRNA.S9469; Nav1; Ocd; olfD; par; sbl; sbl-1; Shu; Shudderer

String
7227.FBpp0303597 (http://string-db.org/newstring_cgi/show_network_section.pl?identifier= 7227.FBpp0303597)

Sequence Similarities
Belongs to the sodium channel (TC 1.A.1.10) family. Para subfamily.

GO - Molecular Function
GO:0005509 : calcium ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005509)
GO:0005244 : voltage-gated ion channel activity (https://www.ebi.ac.uk/QuickGO/term/GO:0005244)
GO:0005248 : voltage-gated sodium channel activity (https://www.ebi.ac.uk/QuickGO/term/GO:0005248)
GO:0005272 : sodium channel activity

UniProtKB Drosophila melanogaster
P35500 (http://www.uniprot.org/uniprot/P35500)

GenebankID or UniProtKB
()

(https://www.ebi.ac.uk/QuickGO/term/GO:0005272)

GO - Biological Process

GO:0045433 : male courtship behavior, veined wing generated song production
(https://www.ebi.ac.uk/QuickGO/term/GO:0045433)
GO:0001666 : response to hypoxia (https://www.ebi.ac.uk/QuickGO/term/GO:0001666)
GO:0009612 : response to mechanical stimulus
(https://www.ebi.ac.uk/QuickGO/term/GO:0009612)
GO:0034765 : regulation of ion transmembrane transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0034765)
GO:0035725 : sodium ion transmembrane transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0035725)
GO:0007638 : mechanosensory behavior
(https://www.ebi.ac.uk/QuickGO/term/GO:0007638)
GO:0060078 : regulation of postsynaptic membrane potential
(https://www.ebi.ac.uk/QuickGO/term/GO:0060078)

GO - Cellular Component

GO:0005887 : integral component of plasma membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:0005887)
GO:0001518 : voltage-gated sodium channel complex
(https://www.ebi.ac.uk/QuickGO/term/GO:0001518)

Presumptive Null

No (https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title)

Molecular Type

Coding (https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title)

Aberration Type

SNP (https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

V1848I

Experimental Evidence

Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Val	Ile	1881

Main Reference

Two novel sodium channel mutations associated with resistance to indoxacarb and metaflumizone in the diamondback moth, Plutella xylostella. (2016)
(https://pubmed.ncbi.nlm.nih.gov/25850422)

Authors

Wang XL; Su W; Zhang JH; Yang YH; Dong K; Wu YD

Abstract

Indoxacarb and metaflumizone belong to a relatively new class of sodium channel blocker insecticides (SCBIs). Due to intensive use of indoxacarb, field-evolved indoxacarb resistance has been reported in several lepidopteran pests, including the diamondback moth Plutella xylostella, a serious pest of cruciferous crops. In particular, the BY12 population of P. xylostella, collected from Baiyun, Guangdong Province of China in 2012, was 750-fold more resistant to indoxacarb and 70-fold more resistant to metaflumizone compared with the susceptible Roth strain. Comparison of complementary DNA sequences encoding the sodium channel genes of Roth and BY12 revealed two point mutations (F1845Y and V1848I) in the sixth segment of domain IV of the PxNav protein in the BY population. Both mutations are located within a highly conserved sequence region that is predicted to be involved in the binding sites of local anesthetics and SCBIs based on mammalian sodium channels. A significant correlation was observed among 10 field-collected populations between the mutant allele (Y1845 or I1848) frequencies (1.7% to 52.5%) and resistance levels to both indoxacarb (34- to 870-fold) and metaflumizone (1- to 70-fold). The two mutations were never found to co-exist in the same allele of PxNav , suggesting that they arose independently. This is the first time that sodium channel mutations have been associated with high levels of resistance to SCBIs. F1845Y and V1848I are molecular markers for resistance monitoring in the diamondback moth and possibly other insect pest species.

Â© 2015 Institute of Zoology, Chinese Academy of Sciences.

Additional References

Functional validation of target-site resistance mutations against sodium channel blocker insecticides (SCBIs) via molecular modeling and genome engineering in Drosophila. (2019)
(https://pubmed.ncbi.nlm.nih.gov/30572019)

RELATED GEPHE

Related Genes

10 (ABCC2, Acetylcholinesterase (Ace-1), Chitin synthase 1 (CHS1), CYP6BG1, FMO2, glutamate-gated chloride channel (GluCl), MAP4K4, nAChR, resistance to dieldrin, RYR)
(https://www.gephebase.org/search-criteria?/or+Taxon ID=^51655^/and+Trait=Xenobiotic resistance/and+groupHaplotypes=true#gephebase-summary-title)

Related Haplotypes

4 (https://www.gephebase.org/search-criteria?/or+Gene Gephebase=^para (kdr)^/and+Taxon ID=^51655^/or+Gene Gephebase=^para (kdr)^/and+Taxon ID=^51655^#gephebase-summary-title)

EXTERNAL LINKS

COMMENTS