GEPHE SUMMARY

Gephebase Gene
Na/K-ATPase alpha-subunit (https://www.gephebase.org/search-criteria?/and+Gene Gephebase=^Na/K-ATPase alpha-subunit^#gephebase-summary-title)

GepheID
GP00000704

Entry Status
Published

Main curator
Courtier

PHENOTYPIC CHANGE

Trait Category
Physiology (https://www.gephebase.org/search-criteria?/and+Trait Category=^Physiology^#gephebase-summary-title)

Trait
Xenobiotic resistance (cardiac glycosides) (https://www.gephebase.org/search-criteria?/and+Trait=^Xenobiotic resistance (cardiac glycosides)^#gephebase-summary-title)

Trait State in Taxon A
Other beetles

Trait State in Taxon B
Chrysochus auratus

Ancestral State
Taxon A

Taxonomic Status
Interspecific (https://www.gephebase.org/search-criteria?/and+Taxonomic Status=^Interspecific^#gephebase-summary-title)

Taxon A

Latin Name
Coleoptera (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Coleoptera^#gephebase-summary-title)

Common Name
beetles

Synonyms
beetles

Rank
order

Lineage
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola

Parent
Holometabola () - (Rank: cohort) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 33392)

NCBI Taxonomy ID
7041 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 7041)

is Taxon A an Infraspecies?
No

Taxon B

Latin Name
Chrysochus auratus (https://www.gephebase.org/search-criteria?/and+Taxon and Synonyms=^Chrysochus auratus^#gephebase-summary-title)

Common Name
dogbane beetle

Synonyms
dogbane beetle

Rank
species

Lineage
cellular organisms; Eukaryota; Opisthokonta; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Holometabola; Coleoptera; Polyphaga; Cucujiformia; Chrysomeloidea; Chrysomelidae; Eumolpinae; Chrysochus

Parent
Chrysochus () - (Rank: genus) (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 131618)

NCBI Taxonomy ID
131619 (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id= 131619)

is Taxon B an Infraspecies?
No

GENOTYPIC CHANGE

Generic Gene Name
Atp1a1

Synonyms
Atpa-1; BC010319

String
10090.ENSMUSP00000039657 (http://string-db.org/newstring_cgi/show_network_section.pl?identifier= 10090.ENSMUSP00000039657)

Sequence Similarities
Belongs to the cation transport ATPase (P-type) (TC 3.A.3) family. Type IIC subfamily.

GO - Molecular Function
GO:0005524 : ATP binding (https://www.ebi.ac.uk/QuickGO/term/GO:0005524)
GO:0043531 : ADP binding (https://www.ebi.ac.uk/QuickGO/term/GO:0043531)
GO:0019901 : protein kinase binding (https://www.ebi.ac.uk/QuickGO/term/GO:0019901)
GO:0043548 : phosphatidylinositol 3-kinase binding (https://www.ebi.ac.uk/QuickGO/term/GO:0043548)
GO:0005391 : sodium:potassium-exchanging ATPase activity (https://www.ebi.ac.uk/QuickGO/term/GO:0005391)

UniProtKB Mus musculus
Q8VDN2 (http://www.uniprot.org/uniprot/Q8VDN2)

GenebankID or UniProtKB
()

GO:0051087 : chaperone binding (https://www.ebi.ac.uk/QuickGO/term/GO:0051087)
GO:0019904 : protein domain specific binding
(https://www.ebi.ac.uk/QuickGO/term/GO:0019904)
GO:0030506 : ankyrin binding (https://www.ebi.ac.uk/QuickGO/term/GO:0030506)
GO:0016791 : phosphatase activity (https://www.ebi.ac.uk/QuickGO/term/GO:0016791)
GO:0030955 : potassium ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0030955)
GO:0031402 : sodium ion binding (https://www.ebi.ac.uk/QuickGO/term/GO:0031402)
GO:1990239 : steroid hormone binding
(https://www.ebi.ac.uk/QuickGO/term/GO:1990239)

GO - Biological Process

GO:0071383 : cellular response to steroid hormone stimulus
(https://www.ebi.ac.uk/QuickGO/term/GO:0071383)
GO:0006813 : potassium ion transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0006813)
GO:0006814 : sodium ion transport (https://www.ebi.ac.uk/QuickGO/term/GO:0006814)
GO:0071260 : cellular response to mechanical stimulus
(https://www.ebi.ac.uk/QuickGO/term/GO:0071260)
GO:0042493 : response to drug (https://www.ebi.ac.uk/QuickGO/term/GO:0042493)
GO:0008217 : regulation of blood pressure
(https://www.ebi.ac.uk/QuickGO/term/GO:0008217)
GO:0015991 : ATP hydrolysis coupled proton transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0015991)
GO:0030007 : cellular potassium ion homeostasis
(https://www.ebi.ac.uk/QuickGO/term/GO:0030007)
GO:0006883 : cellular sodium ion homeostasis
(https://www.ebi.ac.uk/QuickGO/term/GO:0006883)
GO:1990573 : potassium ion import across plasma membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:1990573)
GO:0036376 : sodium ion export across plasma membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:0036376)
GO:0090662 : ATP hydrolysis coupled transmembrane transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0090662)
GO:0060081 : membrane hyperpolarization
(https://www.ebi.ac.uk/QuickGO/term/GO:0060081)
GO:0086009 : membrane repolarization
(https://www.ebi.ac.uk/QuickGO/term/GO:0086009)
GO:0031947 : negative regulation of glucocorticoid biosynthetic process
(https://www.ebi.ac.uk/QuickGO/term/GO:0031947)
GO:0045822 : negative regulation of heart contraction
(https://www.ebi.ac.uk/QuickGO/term/GO:0045822)
GO:0045823 : positive regulation of heart contraction
(https://www.ebi.ac.uk/QuickGO/term/GO:0045823)
GO:0045989 : positive regulation of striated muscle contraction
(https://www.ebi.ac.uk/QuickGO/term/GO:0045989)
GO:0086004 : regulation of cardiac muscle cell contraction
(https://www.ebi.ac.uk/QuickGO/term/GO:0086004)
GO:0002028 : regulation of sodium ion transport
(https://www.ebi.ac.uk/QuickGO/term/GO:0002028)
GO:0002026 : regulation of the force of heart contraction
(https://www.ebi.ac.uk/QuickGO/term/GO:0002026)

GO - Cellular Component

GO:0016021 : integral component of membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:0016021)
GO:0005886 : plasma membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0005886)
GO:0016324 : apical plasma membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:0016324)
GO:0016020 : membrane (https://www.ebi.ac.uk/QuickGO/term/GO:0016020)
GO:0045121 : membrane raft (https://www.ebi.ac.uk/QuickGO/term/GO:0045121)
GO:0005794 : Golgi apparatus (https://www.ebi.ac.uk/QuickGO/term/GO:0005794)
GO:0032991 : protein-containing complex
(https://www.ebi.ac.uk/QuickGO/term/GO:0032991)
GO:0005783 : endoplasmic reticulum
(https://www.ebi.ac.uk/QuickGO/term/GO:0005783)
GO:0005768 : endosome (https://www.ebi.ac.uk/QuickGO/term/GO:0005768)
GO:0016323 : basolateral plasma membrane
(https://www.ebi.ac.uk/QuickGO/term/GO:0016323)
GO:0005901 : caveola (https://www.ebi.ac.uk/QuickGO/term/GO:0005901)
GO:0030315 : T-tubule (https://www.ebi.ac.uk/QuickGO/term/GO:0030315)
GO:0014069 : postsynaptic density (https://www.ebi.ac.uk/QuickGO/term/GO:0014069)
GO:0014704 : intercalated disc (https://www.ebi.ac.uk/QuickGO/term/GO:0014704)
GO:0043209 : myelin sheath (https://www.ebi.ac.uk/QuickGO/term/GO:0043209)
GO:0042383 : sarcolemma (https://www.ebi.ac.uk/QuickGO/term/GO:0042383)
GO:0005890 : sodium:potassium-exchanging ATPase complex
(https://www.ebi.ac.uk/QuickGO/term/GO:0005890)

Mutation #1

Presumptive Null

No (https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title)

Molecular Type

Coding (https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title)

Aberration Type

SNP (https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

Q111L then L111V + N122H +I315L on one recent gene copy

Experimental Evidence

Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Gln	Leu	111

Main Reference

The evolution of cardenolide-resistant forms of Naâº,Kâº -ATPase in Danainae butterflies. (2012) (https://pubmed.ncbi.nlm.nih.gov/22126595)

Authors

Aardema ML; Zhen Y; Andolfatto P

Abstract

Cardenolides are a class of plant secondary compounds that inhibit the proper functioning of the Na(+) , K(+) -ATPase enzyme in susceptible animals. Nonetheless, many insect species are able to sequester cardenolides for their own defence. These include butterflies in the subfamily Danainae (Family: Nymphalidae) such as the monarch (Danaus plexippus). Previous studies demonstrated that monarchs harbour an asparagine (N) to histidine (H) substitution (N122H) in the Î± subunit of Na(+) , K(+) -ATPase (ATPÎ±) that reduces this enzyme's sensitivity to cardenolides. More recently, it has been suggested that at ATPÎ± position 111, monarchs may also harbour a leucine (L)/glutamine (Q) polymorphism. This later amino acid could also contribute to cardenolide insensitivity. However, here we find that incorrect annotation of the initially reported DNA sequence for ATPÎ± has led to several erroneous conclusions. Using a population genetic and phylogenetic analysis of monarchs and their close relatives, we show that an ancient Q111L substitution occurred prior to the radiation of all Danainae, followed by a second substitution at the same site to valine (V), which arose before the diversification of the Danaus genus. In contrast, N122H appears to be a recent substitution specific to monarchs. Surprisingly, examination of a broader insect phylogeny reveals that the same progression of amino acid substitutions (Q111Lâ€ƒâ†'â€ƒL111Vâ€ƒ+â€ƒN122H) has also occurred in Chyrsochus beetles (Family: Chrysomelidae, Subfamily: Eumolpinae) that feed on cardenolide-containing host plants. The parallel pattern of amino acid substitution in these two distantly related lineages is consistent with an adaptive role for these substitutions in reducing cardenolide sensitivity and suggests that their temporal order may be limited by epistatic interactions.

Â© 2011 Blackwell Publishing Ltd.

Additional References

Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. (2012) (https://pubmed.ncbi.nlm.nih.gov/22826239)

Mutation #2

Presumptive Null

No (https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title)

Molecular Type

Coding (https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title)

Aberration Type

SNP (https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

Q111L then L111V + N122H +I315L on one recent gene copy

Experimental Evidence

Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Leu	Val	111

Main Reference

The evolution of cardenolide-resistant forms of Naâº,Kâº -ATPase in Danainae butterflies. (2012) (https://pubmed.ncbi.nlm.nih.gov/22126595)

Authors

Aardema ML; Zhen Y; Andolfatto P

Abstract

Cardenolides are a class of plant secondary compounds that inhibit the proper functioning of the Na(+) , K(+) -ATPase enzyme in susceptible animals. Nonetheless, many insect species are able to sequester cardenolides for their own defence. These include butterflies in the subfamily Danainae (Family: Nymphalidae) such as the monarch (Danaus plexippus). Previous studies demonstrated that monarchs harbour an asparagine (N) to histidine (H) substitution (N122H) in the Î± subunit of Na(+) , K(+) -ATPase (ATPÎ±) that reduces this enzyme's sensitivity to cardenolides. More recently, it has been suggested that at ATPÎ± position 111, monarchs may also harbour a leucine (L)/glutamine (Q) polymorphism. This later amino acid could also contribute to cardenolide insensitivity. However, here we find that incorrect annotation of the initially reported DNA sequence for ATPÎ± has led to several erroneous conclusions. Using a population genetic and phylogenetic analysis of monarchs and their close relatives, we show that an ancient Q111L substitution occurred prior to the radiation of all Danainae, followed by a second substitution at the same site to valine (V), which arose before the diversification of the Danaus genus. In contrast, N122H appears to be a recent substitution specific to monarchs. Surprisingly, examination of a broader insect phylogeny reveals that the same progression of amino acid substitutions (Q111Lâ€ƒâ†'â€ƒL111Vâ€ƒ+â€ƒN122H) has also occurred in Chyrsochus beetles (Family: Chrysomelidae, Subfamily: Eumolpinae) that feed on cardenolide-containing host plants. The parallel pattern of amino acid substitution in these two distantly related lineages is consistent with an adaptive role for these substitutions in reducing cardenolide sensitivity and suggests

that their temporal order may be limited by epistatic interactions.

Â© 2011 Blackwell Publishing Ltd.

Additional References

Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. (2012) (https://pubmed.ncbi.nlm.nih.gov/22826239)

Mutation #3

Presumptive Null

No (https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title)

Molecular Type

Coding (https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title)

Aberration Type

SNP (https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

Q111L then L111V + N122H +I315L on one recent gene copy

Experimental Evidence

Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Asn	His	122

Main Reference

The evolution of cardenolide-resistant forms of Naâº,Kâº -ATPase in Danainae butterflies. (2012) (https://pubmed.ncbi.nlm.nih.gov/22126595)

Authors

Aardema ML; Zhen Y; Andolfatto P

Abstract

Cardenolides are a class of plant secondary compounds that inhibit the proper functioning of the Na(+) , K(+) -ATPase enzyme in susceptible animals. Nonetheless, many insect species are able to sequester cardenolides for their own defence. These include butterflies in the subfamily Danainae (Family: Nymphalidae) such as the monarch (Danaus plexippus). Previous studies demonstrated that monarchs harbour an asparagine (N) to histidine (H) substitution (N122H) in the Î± subunit of Na(+) , K(+) -ATPase (ATPÎ±) that reduces this enzyme's sensitivity to cardenolides. More recently, it has been suggested that at ATPÎ± position 111, monarchs may also harbour a leucine (L)/glutamine (Q) polymorphism. This later amino acid could also contribute to cardenolide insensitivity. However, here we find that incorrect annotation of the initially reported DNA sequence for ATPÎ± has led to several erroneous conclusions. Using a population genetic and phylogenetic analysis of monarchs and their close relatives, we show that an ancient Q111L substitution occurred prior to the radiation of all Danainae, followed by a second substitution at the same site to valine (V), which arose before the diversification of the Danaus genus. In contrast, N122H appears to be a recent substitution specific to monarchs. Surprisingly, examination of a broader insect phylogeny reveals that the same progression of amino acid substitutions (Q111Lâ€ƒâ†'â€ƒL111Vâ€ƒ+â€ƒN122H) has also occurred in Chyrsochus beetles (Family: Chrysomelidae, Subfamily: Eumolpinae) that feed on cardenolide-containing host plants. The parallel pattern of amino acid substitution in these two distantly related lineages is consistent with an adaptive role for these substitutions in reducing cardenolide sensitivity and suggests that their temporal order may be limited by epistatic interactions.

Â© 2011 Blackwell Publishing Ltd.

Additional References

Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. (2012) (https://pubmed.ncbi.nlm.nih.gov/22826239)

Mutation #4

Presumptive Null

No (https://www.gephebase.org/search-criteria?/and+Presumptive Null=^No^#gephebase-summary-title)

Molecular Type

Coding (https://www.gephebase.org/search-criteria?/and+Molecular Type=^Coding^#gephebase-summary-title)

Aberration Type

SNP (https://www.gephebase.org/search-criteria?/and+Aberration Type=^SNP^#gephebase-summary-title)

SNP Coding Change

Nonsynonymous

Molecular Details of the Mutation

Q111L then L111V + N122H +I315L on one recent gene copy

Experimental Evidence

Candidate Gene (https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Candidate Gene^#gephebase-summary-title)

	Taxon A	Taxon B	Position
Codon	-	-	-
Amino-acid	Ile	Leu	315

Main Reference

The evolution of cardenolide-resistant forms of Naâº,Kâº -ATPase in Danainae butterflies. (2012) (https://pubmed.ncbi.nlm.nih.gov/22126595)

Authors

Aardema ML; Zhen Y; Andolfatto P

Abstract

Cardenolides are a class of plant secondary compounds that inhibit the proper functioning of the Na(+) , K(+) -ATPase enzyme in susceptible animals. Nonetheless, many insect species are able to sequester cardenolides for their own defence. These include butterflies in the subfamily Danainae (Family: Nymphalidae) such as the monarch (Danaus plexippus). Previous studies demonstrated that monarchs harbour an asparagine (N) to histidine (H) substitution (N122H) in the Î± subunit of Na(+) , K(+) -ATPase (ATPÎ±) that reduces this enzyme's sensitivity to cardenolides. More recently, it has been suggested that at ATPÎ± position 111, monarchs may also harbour a leucine (L)/glutamine (Q) polymorphism. This later amino acid could also contribute to cardenolide insensitivity. However, here we find that incorrect annotation of the initially reported DNA sequence for ATPÎ± has led to several erroneous conclusions. Using a population genetic and phylogenetic analysis of monarchs and their close relatives, we show that an ancient Q111L substitution occurred prior to the radiation of all Danainae, followed by a second substitution at the same site to valine (V), which arose before the diversification of the Danaus genus. In contrast, N122H appears to be a recent substitution specific to monarchs. Surprisingly, examination of a broader insect phylogeny reveals that the same progression of amino acid substitutions (Q111Lâ€ƒâ†'â€ƒL111Vâ€ƒ+â€ƒN122H) has also occurred in Chyrsochus beetles (Family: Chrysomelidae, Subfamily: Eumolpinae) that feed on cardenolide-containing host plants. The parallel pattern of amino acid substitution in these two distantly related lineages is consistent with an adaptive role for these substitutions in reducing cardenolide sensitivity and suggests that their temporal order may be limited by epistatic interactions.

Â© 2011 Blackwell Publishing Ltd.

Additional References

Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. (2012) (https://pubmed.ncbi.nlm.nih.gov/22826239)

RELATED GEPHE

Related Genes

6 (ABCB1, Acetylcholinesterase (Ace-2), Acetylcholinesterase (Ace), para (kdr), nAChR, resistance to dieldrin) (https://www.gephebase.org/search-criteria?/or+Taxon ID=^7041^/and+Trait=Xenobiotic resistance/or+Taxon ID=^131619^/and+Trait=Xenobiotic resistance/and+groupHaplotypes=true#gephebase-summary-title)

Related Haplotypes

3 (https://www.gephebase.org/search-criteria?/or+Gene Gephebase=^Na/K-ATPase alpha-subunit^/and+Taxon ID=^7041^/or+Gene Gephebase=^Na/K-ATPase alpha-subunit^/and+Taxon ID=^131619^#gephebase-summary-title)

EXTERNAL LINKS

COMMENTS

@SeveralMutationsWithEffect @SuccessiveMutationsAtSameCodon