

## GEPHE SUMMARY

	Gephebase Gene		GepheID
PRR37-like Photoperiod-H1 (Ppd-H1) ( <a href="https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=~PRR37-like+Photoperiod-H1+(Ppd-H1)^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Gene+Gephebase=~PRR37-like+Photoperiod-H1+(Ppd-H1)^#gephebase-summary-title</a> )		GP00000933	
	Entry Status	Martin	Main curator
Published			

## PHENOTYPIC CHANGE

	Trait Category		
Physiology ( <a href="https://www.gephebase.org/search-criteria?/and+Trait+Category=~Physiology^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Trait+Category=~Physiology^#gephebase-summary-title</a> )			
	Trait		
Flowering time ( <a href="https://www.gephebase.org/search-criteria?/and+Trait=~Flowering+time^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Trait=~Flowering+time^#gephebase-summary-title</a> )			
	Trait State in Taxon A		
Hordeum vulgare - (winter)			
	Trait State in Taxon B		
Hordeum vulgare - (spring)			
	Ancestral State		
Data not curated			
	Taxonomic Status		
Domesticated ( <a href="https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=~Domesticated^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Taxonomic+Status=~Domesticated^#gephebase-summary-title</a> )			
Taxon A		Taxon B	
	Latin Name		Latin Name
Hordeum vulgare ( <a href="https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Hordeum+vulgare^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Hordeum+vulgare^#gephebase-summary-title</a> )		Hordeum vulgare ( <a href="https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Hordeum+vulgare^#gephebase-summary-title">https://www.gephebase.org/search-criteria?/and+Taxon+and+Synonyms=~Hordeum+vulgare^#gephebase-summary-title</a> )	
	Common Name		Common Name
-		-	
	Synonyms		Synonyms
barley; Hordeum vulgare L.; Horedum vulgare		barley; Hordeum vulgare L.; Horedum vulgare	
	Rank		Rank
species		species	
	Lineage		Lineage
cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliophyta; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; BOP clade; Pooideae; Triticeae; Triticeae; Hordeinae; Hordeum		cellular organisms; Eukaryota; Viridiplantae; Streptophyta; Streptophytina; Embryophyta; Tracheophyta; Euphyllophyta; Spermatophyta; Magnoliophyta; Mesangiospermae; Liliopsida; Petrosaviidae; commelinids; Poales; Poaceae; BOP clade; Pooideae; Triticeae; Triticeae; Hordeinae; Hordeum	
	Parent		Parent
Hordeum () - (Rank: genus) ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4512">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4512</a> )		Hordeum () - (Rank: genus) ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4512">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4512</a> )	
	NCBI Taxonomy ID		NCBI Taxonomy ID
4513 ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4513">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4513</a> )		4513 ( <a href="https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4513">https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=4513</a> )	
	is Taxon A an Intraspecies?		is Taxon B an Intraspecies?
No		No	

## GENOTYPIC CHANGE

	Generic Gene Name		UniProtKB Oryza sativa subsp. japonica
PRR37		Q0D3B6 ( <a href="http://www.uniprot.org/uniprot/Q0D3B6">http://www.uniprot.org/uniprot/Q0D3B6</a> )	
	Synonyms		GenebankID or UniProtKB
PRR37; OsPRR37; DTH7; HD2; Os07g0695100; LOC_Os07g49460; P0627E10.21		()	
	String		
39947.LOC_Os07g49460.1 ( <a href="http://string-db.org/newstring.cgi/show_network_section.pl?identifier=39947.LOC_Os07g49460.1">http://string-db.org/newstring.cgi/show_network_section.pl?identifier=39947.LOC_Os07g49460.1</a> )			
	Sequence Similarities		
Belongs to the ARR-like family.			
	GO - Molecular Function		
-			
	GO - Biological Process		
GO:0009908 : flower development ( <a href="https://www.ebi.ac.uk/QuickGO/term/GO:0009908">https://www.ebi.ac.uk/QuickGO/term/GO:0009908</a> )			
GO:0000160 : phosphorelay signal transduction system ( <a href="https://www.ebi.ac.uk/QuickGO/term/GO:0000160">https://www.ebi.ac.uk/QuickGO/term/GO:0000160</a> )			
GO:0009585 : red, far-red light phototransduction			

(<https://www.ebi.ac.uk/QuickGO/term/GO:0009585>)  
GO:0048579 : negative regulation of long-day photoperiodism, flowering  
(<https://www.ebi.ac.uk/QuickGO/term/GO:0048579>)  
GO:0048511 : rhythmic process (<https://www.ebi.ac.uk/QuickGO/term/GO:0048511>)  
GO - Cellular Component  
GO:0005634 : nucleus (<https://www.ebi.ac.uk/QuickGO/term/GO:0005634>)

Presumptive Null

Unknown (<https://www.gephebase.org/search-criteria?/and+Presumptive Null=^Unknown^#gephebase-summary-title>)

Molecular Type

Unknown (<https://www.gephebase.org/search-criteria?/and+Molecular Type=^Unknown^#gephebase-summary-title>)

Aberration Type

Unknown (<https://www.gephebase.org/search-criteria?/and+Aberration Type=^Unknown^#gephebase-summary-title>)

Molecular Details of the Mutation

unknown

Experimental Evidence

Linkage Mapping (<https://www.gephebase.org/search-criteria?/and+Experimental Evidence=^Linkage Mapping^#gephebase-summary-title>)

Main Reference

The pseudo-response regulator Ppd-H1 provides adaptation to photoperiod in barley. (2005) (<https://pubmed.ncbi.nlm.nih.gov/16284181>)

Authors

Turner A; Beales J; Faure S; Dunford RP; Laurie DA

Abstract

Plants commonly use photoperiod (day length) to control the timing of flowering during the year, and variation in photoperiod response has been selected in many crops to provide adaptation to different environments and farming practices. Positional cloning identified Ppd-H1, the major determinant of barley photoperiod response, as a pseudo-response regulator, a class of genes involved in circadian clock function. Reduced photoperiod responsiveness of the ppd-H1 mutant, which is highly advantageous in spring-sown varieties, is explained by altered circadian expression of the photoperiod pathway gene CONSTANS and reduced expression of its downstream target, FT, a key regulator of flowering.

Additional References

Expression conservation within the circadian clock of a monocot: natural variation at barley Ppd-H1 affects circadian expression of flowering time genes, but not clock orthologs. (2012) (<https://pubmed.ncbi.nlm.nih.gov/22720803>)

## RELATED GEPHE

Related Genes

6 (CENTRORADIALIS (HvCEN), EARLY FLOWERING 3 (here = Mat-a), EARLY FLOWERING 3/ EARLY MATURITY 8, EARLY FLOWERING 3/ EARLYMATURITY8, Flowering locus T (=HvFT=VRN3), VRN2) (<https://www.gephebase.org/search-criteria?/or+Taxon ID=^4513^/and+Trait=Flowering time/and+groupHaplotypes=true#gephebase-summary-title>)

Related Haplotypes

No matches found.

## EXTERNAL LINKS

## COMMENTS

@GxE